Geophysical Research Abstracts Vol. 19, EGU2017-6474, 2017 EGU General Assembly 2017 © Author(s) 2017. CC Attribution 3.0 License.



Comparison between OMI-TOMS and OMI-DOAS Ozone Columns over High SO₂ regions: Volcanic and Chinese Industrial Region

Wonei Choi, Hyunkee Hong, and Hanlim Lee

Division of Earth Environmental System Science, Major of Spatial Information Engineering, Pukyong National University, Busan, Korea, Republic Of (cwyh3338@gmail.com)

In this present study, we identified the SO₂ effect on O₃ retrieval from the Ozone Monitoring Instrument (OMI) measurement. The difference between OMI-Total Ozone Mapping Spectrometer (TOMS) and OMI-Differential Optical Absorption Spectrometer (DOAS) total O₃ is calculated in high SO₂ volcanic regions (Anatahan, La Cumbre, Sierra Negra, and Piton) and Chinese industrial region from 2005 through 2008. In volcanic plumes, OMI-TOMS total O₃ column increases with middle and upper troposphere and stratosphere (TRM-STL) SO₂, showing high correlation (R = 0.54) between TRM-STL SO₂ and TOMS O₃ while there is negligible correlation (R = 0.11) between TRM-STL SO₂ and OMI- DOAS O₃. There is a certain correlation (R = 0.54) between TRM-STL SO₂ and the difference between OMI-TOMS and OMI-DOAS O₃ (T-D). The T-D reaches its maximum up to 90 DU. T-D has a dependence on planetary boundary layer (PBL) SO₂ (R = 0.36) even though T-D in the Chinese industrial region increases more slowly with SO₂ than at the volcanic sites. Thus, the altitude at which SO₂ exists has an effect on T-D, which could be due to reduced absolute radiance sensitivity in the boundary layer at 317.5 nm (the wavelength used to retrieve OMI-TOMS ozone in boundary layer). In addition, T-D shows a seasonal tendency on PBL SO₂. In Turkey and Western USA, where PBL SO₂ concentrations are low, |T-D| shows its positive maximum in summer and negative maximum in winter. However, no seasonal variation of T-D was found in the Chinese industrial region due to the effect of low and high PBL SO₂ in summer and winter, respectively. The reason for the seasonal characteristics of T-D in the Chinese industrial region will be discussed.