Geophysical Research Abstracts Vol. 20, EGU2018-11600-1, 2018 EGU General Assembly 2018 © Author(s) 2018. CC Attribution 4.0 license.



Atmospheric gravity waves generated by pulsed ionospheric convection: A link between the solar wind and the Earth's atmosphere

Paul Prikryl (1,2), Robert Bruntz (3), and Bharat S. R. Kunduri (4)

(1) University of New Brunswick, Physics Department, Fredericton, NB, Canada (paul.prikryl@unb.ca), (2) Geomagnetic Laboratory, Natural Resources Canada, Ottawa, ON, Canada, (3) Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA, (4) Bradley Dept. of Electrical and Computer Engineering, Virginia Tech, Blacksburg, VA, USA

Atmospheric gravity waves (AGWs) [1] play an important role in transporting momentum and energy through the atmosphere. Solar wind coupling to the magnetosphere-ionosphere-atmosphere (MIA) system has been recognized as an important source of AGWs that are manifested in the ionosphere as traveling ionospheric disturbances (TIDs) and observed by HF radars [2] and ionosondes. Solar wind Alfvén waves modulating Joule heating and/or Lorentz forcing of the high-latitude lower thermosphere have been shown to generate medium-scale AGWs [3]. Using the dispersion relation between the gravity wave frequency and the wave vector [1], ray tracing [3] in a model atmosphere shows three distinct group paths (wave propagation modes) that reach the F region of the ionosphere: waves that travel directly upward, waves that are reflected in the mesosphere, and waves that are reflected in the troposphere or from the Earth's surface. Another approach is the Transfer Function Model (TFM) [4] that describes gravity wave response in the atmosphere and shows that propagating waves originating in the thermosphere can excite a spectrum of AGWs in the lower atmosphere, albeit with much smaller amplitudes. These include "lower modes" that are of particular interest in the present paper: the wave reflected from the Earth's surface and the ducted wave that propagates in the lower atmosphere to long distances away from the source and leaks up into the thermosphere. We will present observations of TIDs and discuss a possible influence of the downward propagating gravity waves on tropospheric weather [5].

References

- [1] Hines C.O., Can. J. Phys. 38, 1441–1481, 1960.
- [2] Samson J. C., et al., Geophys. Res. Lett., 16, 875–878, 1989.
- [3] Prikryl P., et al., Ann. Geophys., 23, 401-417, 2005, doi:10.5194/angeo-23-401-2005.
- [4] Mayr H.G., et al., J. Geophys. Res., 89, 10929–10959, 1984.
- [5] Prikryl P., et al., J. Atmos. Sol.-Terr. Phys., in press, https://doi.org/10.1016/j.jastp.2017.07.023