



Nitrogen balance in acid sulfate soil lysimeters under different water management

Seija Virtanen (1), Asko Simojoki (2), Kenedy E. Epie (3), and Helinä Hartikainen (4)

(1) Drainage Foundation sr, Helsinki, Finland (seija.virtanen@tukisaatio.fi), (2) University of Helsinki, Helsinki, Finland (asko.simojoki@helsinki.fi), (3) Montana State University-WTARC Conrad, Montana State, USA (etone.kenedy@montana.edu), (4) University of Helsinki, Helsinki, Finland (helina.hartikainen@helsinki.fi)

Reclamation of permanently saturated potential acid sulphate (AS) soils for cultivation or other purposes has negative environmental impacts, as drainage triggers soil acidification that subsequently causes off-site harm, e.g. deterioration of ecosystems in recipient watercourses. In addition, the subsoil layers of AS soils are typically high in N and OC, which gives rise to concern for their potential to produce GHG emissions to atmosphere and high N loads to watercourses.

One option to manage soil acidification is to raise groundwater level. In this study, we investigated the effect of raised water table on the N balance of AS soils in ten meso-scale monolithic lysimeters equipped with water level control and probes e.g. for redox measurements. Reed canary grass (RCG) was cultivated in eight lysimeters: four with a high water table (HWT), and four with a low water table (LWT) mimicking normal drainage of cultivated fields. In spring, the lysimeters were fertilized as recommended for RCG (N 90, P 5 and K 40 kg/ha). They were watered with artificial rain water mimicking the local long-term average precipitation. The discharge water was analyzed for dissolved N (Shimadzu TOC-VCPH/CPN). The total N in the dried plant samples was determined by the Dumas combustion method (VarioMax-CN). Gas emissions were measured by using closed chambers and analysing the gas composition by GC.

The N mass balance was calculated as follows: $N_{balance} = N_{input} - N_{output}$ where the only input term was $N_{fertilization}$ and output terms were $N_{harvest}$, $N_{leaching}$ and $N_{emission}$. In all treatments, the $N_{balance}$ was negative. Even though the RCG dry-matter yield was highest in HWT (16 t/ha vs. 12 t/ha in LWT), the $N_{harvest}$ was highest in LWT (142.2 kg-N/ha vs. 112.7 kg-N/ha in HWT). Furthermore, $N_{leaching}$ (3.6 kg-N/ha in HWT and 2.3 kg-N/ha in LWT) and $N_{emission}$ (cumulative N_2O emission of 4.0 kg-N/ha in 161 days, n.s. difference between HWT and LWT) were lower than reported for AS soils in some field scale studies. The lower $N_{leaching}$ was partly due to the high transpiration and N uptake by RCG. Additionally, the negative $N_{balances}$ obtained were supposedly attributable to large plant-available N pools in the subsoil.

Actually, the N_{min} content in the 15-cm subsoil horizons (C1) in the lysimeters corresponded to 124 kg- N_{min} /ha in the field scale. Evidently, the high $N_{harvest}$ in LWT was attributable to the roots penetrating down to the bottom of lysimeters. In HWT, on the contrary, the roots accumulated mainly in the topsoil. Consequently, the $N_{harvest}$ in HWT was lower than in LWT.

In AS soil fields, low groundwater table maintains continuous soil acidification exerting harmful off-site effects. High water tables facilitates only the growth of water tolerant plants, such as RCG with negative $N_{balance}$. However, high groundwater levels in AS soil may lead to mobilization of iron from secondary minerals (e.g. jarosite) as a response to lowered redox potential in watersaturated soil. In addition, in case the plants are not harvested, the mobilization of N in the subsequent years will be high.