



## **Hyperpectral infrared imaging of volcanic plume at Mt Etna during IMAGETNA campaign**

Charlotte Segonne (1), Nathalie Huret (1), Sébastien Payan (2), Valéry Catoire (1), Giuseppe Salerno (3), Tjarda Roberts (1), Alessandro La Spina (3), and Tommaso Caltabiano (3)

(1) Orléans University and LPC2E/CNRS and, 3A avenue de la recherche scientifique, F4500 Orléans (charlotte.segonne@cnrs-orleans.fr), (2) LATMOS, CNRS, Sorbonne Université, France, (3) Istituto Nazionale di Geofisica e Vulcanologia, Osservatorio Etno, Catania, Italy

Knowledge of the composition and the spatial evolution of volcanic plumes provides insights to processes occurring in the Earth's interior. On the other hand, quantification of gaseous emission fluxes is also a fundamental task in the framework of climate change in order to refine the contribution of natural emissions. UV cameras allow us to image volcanic plumes and evaluate  $\text{SO}_2$  fluxes, although can be subject to uncertainties in the retrieval. Another technique of imaging is now available in the infra-red. Such infrared hyperspectral imager (pixel-by-pixel spectra) might represent a major step forward in volcanology due to its potential to allow  $\text{SO}_2$  flux measurements during the night and gives access to additional relevant species but has to be tested and validated as a first step. In June 2015 a campaign of measurements - IMAGETNA – was performed at Mt Etna (Pizzi Deneri Volcano Observatory) with the intent to explore the application of these techniques for volcanic gas measurements all together. Over five days the volcanic plume was remotely observed simultaneously by employing three different hyperspectral imagers (commercial and under development), FTIR instrument, UV LWIR cameras, and radiometer. Results gathered from IR hyperspectral imager will be presented and compared to  $\text{SO}_2$  UV-Vis camera measurements.