

## Conditions of ductile low-angle normal faulting: The Gurktal extensional allochthon

Oliver Stauber (1), Franz Neubauer (1), Manfred Bernroider (1), Shuyun Cao (1,2), Johann Genser (1), Maurizio Musso (3), and Daniela Reiff (3)

(1) Department Geography and Geology, Salzburg University, Salzburg, Austria (oliver.stauber@sbg.ac.at), (2) State Key Laboratory of Geological Processes and Mineral Resources, School of Earth Sciences, China University of Geosciences, Wuhan, China, (3) Chemistry and Physics of Materials, Salzburg University, Salzburg, Austria

Orogenic thrust wedges are sometimes superposed by large-scale extensional deformation as the result of orogenic collapse, when rheologically weak layers are activated as ductile normal faults during exhumation of higher-grade metamorphic units. A heavily disputed example within the Austroalpine nappe stack is the Gurktal nappe complex, which is underlain by the Bundschuh nappe complex (retrogressed polymetamorphic Bundschuh basement overlain by the monometamorphic Stangalm Mesozoic). Previous estimates of metamorphic conditions indicate that approx. 200 °C of the metamorphic profile is missing between the Bundschuh nappe (approx. 480 - 500 °C) and the higher part (250 - 280 °C) of the Gurktal nappe complex suggesting a metamorphic break in between. To solve this paradox, we studied micro- and meso-scale fabrics of these units supplemented by studies of metamorphic T conditions and Ar-Ar white mica dating. This resulted in new data presented here from footwall to hangingwall units:

Structural evidence indicates a ESE-directed ductile low-angle normal faulting passing from ductile, through ductile-brittle to purely brittle conditions under the same external kinematics. Dolomite microfabrics of the metamorphic Stangalm Mesozoic comprise symmetric and asymmetric fabrics indicating a mixture of coaxial and non-coaxial deformation regimes implying partitioning of shear strain. Calcite-dolomite thermometry gives a bimodal distribution of temperatures with maxima at approx. 360°C and approx. 450°C, while a white mica concentrate gives a plateau age of  $96.2 \pm 0.4$  Ma.

The overlying phyllites and phyllonites of the Phyllonite zone are regionally considered to be part of the Murau nappe of the Gurktal nappe complex. The structure shows fabrics passing from initial ductile to subsequently cataclastic and brittle conditions, illustrating the passage through the ductile-brittle boundary during cooling. Raman spectroscopy indicates a preliminary deformation T of 319 – 382 °C (all Raman calibrations are after Kouketsu et al. 2014, Island Arc 23, 33–50). A white mica concentrate gives a plateau age of  $87.1 \pm 0.5$  Ma (age of ductile shearing), whereas another sample gives a mixed age of  $204.3 \pm 0.5$  Ma suggesting the polymetamorphic nature of the Phyllonite zone.

The higher part of the Gurktal nappe complex is comprised of two structural units, the lower Pfannock basement-cover unit and the overlying Stolzalpe nappe, with a nappe contact in between. Graphitic carbon from a semi-ductile shear zone within Upper Carboniferous conglomerate gives a deformation T between 260 and 288 °C, indicating the temperature of thrusting of the Stolzalpe nappe over the Pfannock basement-cover unit, potentially towards WNW according to preliminary kinematic data, predating the ductile low-angle normal fault at the base of the Pfannock unit.

Consequently, we argue that a nappe stack was formed during late Early Cretaceous continental collision subsequently overprinted by an orogenic collapse stage, which also led to exhumation of high-pressure metamorphic units elsewhere within the Austroalpine nappe stack.