Geophysical Research Abstracts Vol. 20, EGU2018-1409, 2018 EGU General Assembly 2018 © Author(s) 2017. CC Attribution 4.0 license.



Geological \mathbf{CO}_2 storage in saline carbonate formations using core flooding experiment

Shachi Shachi (1), Brijesh Kumar Yadav (1), and Aziz Rahman (2)

(1) Indian Institute of Technology Roorkee, Department of Hydrology, India (shachi2809@gmail.com), (2) Texas A&M University, Qatar, Petroleum Engineering Program (aziz.rahman@qatar.tamu.edu)

Continuous anthropogenic emission of green-house gases in atmosphere resulted in global warming which urgently requires mitigation techniques such as carbon capture and its subsequent sequestration. Geological sequestration of CO₂ in deep saline formations is an emerging technique to reduce carbon concentration in atmosphere. Investigations on CO2 sequestration and its subsequent evolution in subsurface under targeted geological sites are at nascent stage. During the course of CO₂ geo-sequestration process the receiving geological media is exposed to cyclic CO₂-brine flooding. The successful implementation of CO₂ geo sequestration process requires the investigation of the target geological formation for its capacity, integrity, interaction between rock and CO₂/brine system. The objective of this study is to investigate the effect of cyclic CO₂-brine flooding on multiphase flow behavior during geo-sequestration on carbonate core under in-situ geological reservoir conditions. A series of practical experiments are conducted at two salinities 3 % and 7.5 % having 8 and 10 MPa as injection pressure using core flooding apparatus at TAMUQ. The pressure drop across the core increases with salinity and successive injection cycles; and decreases with injection pressure which significantly leads to dynamic response of core samples. Such changes could be due to hysteresis in CO₂ and brine flow through the core samples, formation changes, migration of fines, chemical reactions between carbonate and brine/CO₂. The findings of this study can be utilized in commencing any CO₂ geo-sequestration study in deep saline carbonate subsurface to mitigate the impact of greenhouse gas emissions.