

## Observations of mixed warm and cold electrons with RPC-MIP at comet 67P/Churyumov-Gerasimenko

Nicolas Gilet (1), Pierre Henri (1), Anders Eriksson (2), Xavier Vallières (1), Gaëtan Wattieaux (3), Ingo Richter (4), Jérôme Moré (1), Orélien Randriamboarison (1), Elias Odelstad (2,5), Fredrik Johansson (2,5)

(1) Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS, Université d'Orléans, Orléans, France (ngilet@cnrs-orleans.fr), (2) Swedish Institute of Space Physics, Uppsala, Sweden, (3) Université de Toulouse, LAPLACE-UMR 5213, Toulouse, France, (4) Institut für Geophysik und extraterrestrische Physik, TU Braunschweig, Braunschweig, Germany, (5) Department of Physics and Astronomy, Uppsala University, Sweden

In the ionosphere of the comet 67P/Churyumov-Gerasimenko, the Rosetta Plasma Consortium (RPC) instruments have been measuring the plasma environment during more than two years from August 2014 to September 2016. The Ion Electron Sensor (RPC-IES) and the Langmuir Probes (RPC-LAP) onboard the Rosetta orbiter reported different electron populations: (i) a suprathermal electron component (40-100 eV), (ii) a warm electron component of cometary origin (5-10 eV) and (iii) a cold component (<1 eV) of electrons thermalized by collisions with neutrals.

The Mutual Impedance Probe (RPC-MIP) has been operated to measure the total electron density. Gilet et al. (2017) simulated the mutual impedance response for a probe immersed in a two-electron temperature plasma, with both cold and warm component modelled by a Maxwellian distribution.

Through a direct comparison between simulated and measured mutual impedance spectra, the density and the temperature of the two electron populations have been retrieved. This study focuses on three events: (i) on 2015 November 1st in the magnetized cometary plasma near perihelion (1.4 AU), (ii) inside a diamagnetic (unmagnetized) region on 2015 November 20, still near perihelion, and (iii) far from perihelion (3.2 AU) on 2016 May 23.

We illustrate the dynamics of the cold and warm electron populations and show that cold electrons can be transported along the magnetic field far from the electron-neutral collision dominated region where electrons are expected to have cooled down.