



High-resolution late Holocene record from lake Bayan Nuur (Uvs Aimag, Mongolia): the multi-proxy study

Natalia Rudaya (1,2,3), Larisa Frolova (4), Olga Palagushkina (4), Larisa Nazarova (2,4), Demberel Otgonbayar (6), and Liudmila Syrykh (5)

(1) Institute of Archaeology and Ethnography SB RAS, Centre of Cenozoic Geochronology, Novosibirsk, Russia, (2) Potsdam University, Potsdam, Germany, (3) AWI Potsdam, Germany, (4) Kazan Federal University, Kazan, Russia, (6) Khovd State University, Khovd, Mongolia, (5) Herzen State Pedagogical University of Russia

Freshwater Lake Bayan Nuur (49.98 N, 93.95 E, 932 m a.s.l.) is located in the north of the Bor Khyariin Els sand dune in Zuun Govi Soum, Uvs aimag. It was previously investigated by Grunert et al. (2000); and then Tian et al. (2014) done a quantitative reconstruction of climate based on the pollen data for late glacial–Holocene. The time resolution of core sampling has not been very representative (210 yr). We present the new study with high-time resolution (ca 39 yr) covered last 3.8 kyr. The core BN2016 was obtained in 2016 from the depth of 30 m and was subdivided into several sets of samples for 14C; TOC and 13C/12C; diatom, cladocera and chironomid analyses. The funding of the work of NR has been provided by the Alexander von Humboldt Foundation (Ref 3.3 - RUS - 1151158 - HFST-E).

References:

Grunert J. et al., 2000 *Quaternary International* 65/66, 171-192. Tian F. et al., 2014 *Journal of Biogeography*, 1-14.