



## **Grazing on mycorrhizae by the rhizosphere food web regulates seedling survival**

Sina Adl

University of Saskatchewan, Agriculture and Bioresources, Soil Sciences, Saskatoon, Canada (sina.adl@usask.ca)

We selected the eastern white pine (*Pinus strobus*) germinated in soil microcosms, in a full factorial design with 5 treatments each randomly replicated in ten trays (n=10, 50 seedlings) in a controlled-environment greenhouse. The pine is an obligate mycorrhizal species, and the five treatments provided more or less fungivory on the mycorrhizal association during the first 120 days of growth. Treatments were a) no ectomycorrhizae; b) 3 species of ectomycorrhizae (EM); c) EM and enriched with fungivorous nematode culture (Nf); d) EM, Nf, and enriched with microarthropods (MA); e) EM and MA. Seedlings were destructively sampled at four months or at seedling death (<4 months). Data were analysed with ANOVA and multivariate procedures and results were compared across treatments for seedling survival rate, age at death, growth, mycorrhization and animal abundance and composition. Canonical scores plot in discriminant analysis clearly separated treatments by mean (%) root tip colonization. We observed, 1) a trophic cascade, confirming an earlier microcosm study by the Setälä laboratory, and expanding the results to germinating seedlings during mycorrhization; 2) that fungivory in the rhizosphere can decrease mean mycorrhization levels, and affects both seedling survival and growth.

Next we used stable isotopes tracer studies to measure the rate of fungivory by three microarthropods and their predator, for nutrient transfer through trophic levels. Excretion rates were also determined as well as moulting rate for the collembolan *Lepidocyrtus*, and food processing efficiency for the mite (Astigmata) *Tyrophagus*, the mites (Oribatida) *Oribatula* sp, *Hypoaspis* sp and the predatory mite (Mesostigmata) *Lepidocyrtus* sp. We also obtained functional response curves of three fungivorous nematodes on the mycorrhizae.

Our results show that we can measure stable isotope enrichment for single microarthropods using our modified mass spectrometry technique, that fungivory rates can be measured, and confirm our microcosm results, that grazing pressure on early stages of mycorrhization reduces seedling growth and can lead to seedling mortality.