



## **Evaluation and adjustment of description of denitrification in the DailyDayCentandCoup models based on N<sub>2</sub> and N<sub>2</sub>O laboratory incubation system measurements**

Balazs Grosz (1), Reinhard Well (1), Jan-Reent Köster (2), Simone Merl (2), and Bianca Ziehmer (2)

(1) Thünen Institute, Climate-Smart Agriculture, Braunschweig, Germany (reinhard.well@thuenen.de), (2) Ministerium für Umwelt, Energie, Ernährung und Forsten von Rheinland-Pfalz, Mainz, Germany

Denitrification is an anaerobic key process by microbes where the NO<sub>3</sub><sup>-</sup> is step-by-step reduced and emitted as NO, N<sub>2</sub>O and finally N<sub>2</sub> gas from the soil. The accurate knowledge of the reduction of nitrate (NO<sub>3</sub><sup>-</sup>) and nitrite (NO<sub>2</sub><sup>-</sup>) to N<sub>2</sub>O and molecular N<sub>2</sub> is important because the N<sub>2</sub>O fraction is further reduced to N<sub>2</sub> and constitutes the main emission source of this greenhouse gas from agricultural soils. Hence, our understanding and ability to quantify soil denitrification is crucial for mitigating nitrogen fertilizer loss as well as for reducing N<sub>2</sub>O emissions. Models can be an important tool to predict mitigation effects and help to develop climate-smart mitigation strategies.

Ideally, commonly used biogeochemical models could describe the main processes and provide adequate predictions of denitrification processes of agricultural soils but often simplified process descriptions and inadequate model parameters prevent models from simulating adequate fluxes of N<sub>2</sub> and N<sub>2</sub>O on field scale. Model development and parametrization often suffer from limited availability of empirical data describing denitrification processes in agricultural soils. While in many studies N<sub>2</sub>O emissions are used to develop and train models, detailed measurements on NO, N<sub>2</sub>O, N<sub>2</sub> fluxes and concentrations and related soil conditions are necessary to develop and test adequate model algorithms. Composition of denitrifying communities, coinciding effects of management and local conditions on the development of denitrification hotspots are highly variable in space and time.

To address this issue the coordinated research unit „Denitrification in Agricultural Soils: Integrated Control and Modelling at Various Scales (DASIM)” was initiated to investigate more closely N-fluxes caused by denitrification in response to environmental effects, soil properties and microbial communities.

Data suitable to validate denitrification models are still scarce due to previous technical and/or methodical limitations of measuring N<sub>2</sub> fluxes, but large data-sets are needed in view of the extreme spatio-temporal heterogeneity of denitrification. We used two datasets for the model evaluation. The results of the DASIM project and an older but similar experiment will provide such data base on laboratory incubations including measurement of N<sub>2</sub>O and N<sub>2</sub> fluxes and determination of the relevant drivers.

Here, we present how we will use these data to evaluate common biogeochemical process models (CoupModel and the nitrification and denitrification sub-models of DailyDayCent) with respect to modeled N<sub>2</sub>O and N<sub>2</sub> fluxes from denitrification. Using measured input and output parameters will provide a better understanding of the limitations of the tested models and thus be a basis for future model improvement.