

Water during core/mantle segregation: implications on hydrogen incorporation in the core of the Earth and Mars

Vincent Clesi (1), Mohamed Ali Bouhifd (1), Nathalie Bolfan-Casanova (1), Geeth Manthilake (1), Federica Schiavi (1), Caroline Raepsaet (2), Hélène Bureau (2), Hicham Khodja (3), and Denis Andrault (1)

(1) Laboratoire Magmas et Volcans, Aubière, France (v.clesi@opgc.univ-bpclermont.fr), (2) Laboratoire d'Etudes des Elements Légers CEA, Gif-Sur-Yvette, France, (3) Institut de Minéralogie et de Physique de la Matière et de Cosmochimie, Paris, France

The accretion of planets from primordial materials and their subsequent differentiation to form a core and a mantle are fundamental questions in terrestrial and solar system [1-3]. Many of the questions about the processes are still open and much debated. For example, could the presence of water during the metallic phase segregation affect the planet-accretion models? The existing studies on the elemental metal-silicate partitioning under hydrous conditions are limited to conditions not representative of those prevailing during core formation on Earth and Mars [e.g. 4,5]. In the present study we use multi-anvil technique over a large range of P, T, fO₂ and water content.

The present experimental results show that except for Fe, there is no effect of water on the partitioning of moderately siderophile elements. It allowed us to build consistent models of planetary accretion from reducing to oxidized conditions. Furthermore, for the range of water concentrations studied, there is no evidence of an important hydrogen incorporation into planetary cores, thus making unlikely for hydrogen to be a major light element of the core as previously assumed [4,5].

[1] Morbidelli et al. (2000) Meteoritics & Planetary Science 1320, 1309-1320. [2] Marty (2012) EPSL 313-314, 56-66. [3] Sarafian et al. (2014) Science 346, 623-626. [4] Righter and Drake (1999) EPSL 171, 383-399. [5] Okuchi, T. (1997) Science 278, 1781-1784