



The salience of nonlinearities in the boreal winter response to ENSO

Chaim Garfinkel (1), Israel Weinberger (1), Luke Oman (2), Valentina Aquila (3), Ian White (1), and Young-Kwon Lim (2)

(1) Hebrew University of Jerusalem, Earth Science Institute, Jerusalem, Israel (chaim.garfinkel@mail.huji.ac.il), (2) NASA GSFC, (3) American University

A series of simulations using the NASA Goddard Earth Observing System Chemistry-Climate Model are analyzed in order to assess the prominence of nonlinearities in the response to the El Niño - Southern Oscillation (ENSO). In the Central North Pacific region where the sea level pressure response to ENSO peaks, nonlinearities are relatively muted. In contrast, changes to the east of this region (i.e. the far-Northeastern Pacific) and to the north of this region (over Alaska) in response to different ENSO phases are more clearly nonlinear, and become statistically robust after more than 15 events are considered. The relative prominence of these nonlinearities is related to the zonal wavenumber of the tropical precipitation response. Associated with these nonlinearities over the far-Northeastern Pacific are nonlinearities in precipitation over Southwestern United States and surface temperature over Northwest North America and Midwestern United States. In all regions at least 15 events of each type are necessary before nonlinearities can be identified as statistically significant at the 95% confidence level, and as there have only been a similar number of ENSO event of each type since 1920, it is not surprising that it has been difficult to establish statistically significant nonlinearities using observational data.