

Impacts of polar ice sheets on the East Asian monsoon during the MIS-13 interglacial

Feng Shi (1,2), Qiuzhen Yin (1), Irina Nikolova (1,3), Sen Zhao (4,5), Yong Sun (6,7), Gilles Ramstein (7), Zhengtang Guo (2,8,9), and André Berger (1)

(1) Georges Lemaître Centre for Earth and Climate Research, Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve 1348, Belgium , (2) Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China, (3) School of Geography, Earth & Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, UK, (4) Key Laboratory of Meteorological Disaster of Ministry of Education, and College of Atmospheric Science, Nanjing University of Information Science and Technology, Nanjing, 210044, China, (5) School of Ocean and Earth Sciences and Technology, University of Hawaii at Manoa, Honolulu, HI 96822, USA , (6) State Key Laboratory of Numerical Modelling for Atmospheric Sciences and Geophysical Fluid Dynamis (LASG), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China, (7) Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette F-91191, France, (8) CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China, (9) University of Chinese Academy of Sciences, Beijing 100049, China

Among all the interglacials of the last one million years, Marine Isotope Stage (MIS) 13 has the highest $\delta^{18}\text{O}$ value over the past 800 ka in the deep-sea sediments. This would indicate that MIS-13 is the coolest interglacial if assuming $\delta^{18}\text{O}$ mainly represents global ice volume. The Antarctic ice core records show also that MIS-13 is the coolest interglacial over Antarctica with almost the lowest greenhouse gases concentrations (GHG). However, many proxy records from the northern hemisphere (NH) indicate that MIS-13 is at least as warm as or even warmer than the recent interglacials, with extremely strong summer monsoon and a possible melting of Greenland ice sheet. In this study, based on proxy reconstructions, different scenarios regarding the size of the Greenland and Antarctic ice sheets are made, and the response of the East Asian summer monsoon to these scenarios are tested by using the HadCM3 model as well as factor separation analysis and under the astronomical and GHG configurations of MIS-13. The results show that the influence of the disappearance of Greenland ice sheet on the surface temperature is quite local, mainly over the northern high latitudinal regions. However, the influence of the bigger Southern Hemisphere ice sheet on the surface temperature is very global, but especially in the southern hemisphere. This ice sheet condition has an impact on the precipitation pattern over tropical-subtropical regions. It increases summer precipitation over all the East Asian monsoon region, in line with proxy reconstructions from China. The mechanism is related to the thermodynamic and topographical effects of the Greenland and Antarctic ice sheets that influence the East Asian monsoon through atmosphere teleconnections. The scenario of melted Greenland ice sheet and of larger SH ice sheets contributes to the explanations of a strong monsoon rainfall in China during MIS-13 as documented by proxy data.