

Origin and historical inputs of suspended particulate matter (SPM) from the Rhône tributaries: use of the non-reactive geochemical signature of particles.

Céline Bégorre ⁽¹⁾, Aymeric Dabrin ⁽¹⁾, Matthieu Masson ⁽¹⁾, Ghislaine Grisot ⁽¹⁾, Lysiane Dherret ⁽¹⁾, Frédérique Eyrolle-Boyer ⁽²⁾, Alain Veron ⁽³⁾, Brice Mourier ⁽⁴⁾, and Marina Coquery ⁽¹⁾

Contact: celine.begorre@irstea.fr

(1) Irstea, UR RiverLy, centre de Lyon-Villeurbanne, 5 rue de la Doua CS 20244, 69625 Villeurbanne, France, (2) Institut de Radioprotection et de Sureté Nucléaire (IRSN), PSE-ENV, SRTE/LRTA, BP 3, Saint-Paul-lez-Durance, France, (3) CEREGE, Aix-Marseille Univ., CNRS, IRD, Coll de France, Europôle Méditerranéen de l'Arbois, BP80, Aix en Provence Cedex 04, France, (4) Université de Lyon, UMR5023 Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, Université Lyon 1, ENTPE, CNRS, 3, rue Maurice Audin, 69518 Vaulx-en-Velin, France

Vienna, 2018



Context


- The determination of sediment origin is essential to manage suspended particulate matter inputs in river systems.
- SPM contribute to the transport of a large amount of contaminants in rivers.
- The Rhône River is the main sediment input to the Mediterranean Sea. The Rhône sediment observatory (OSR) program aims to develop a fingerprinting approach in Rhône River watershed with a large number of SPM samples over 7 years (n = 300).

Objectives


- Determine the origin of SPM from the Rhône River tributaries :
 - Estimate relative contributions of SPM fluxes using conservative tracers (major and trace elements in the residual fraction, particle size correction)
 - Estimate uncertainties associated with these contributions
- Determine the historical SPM inputs of the tributaries by applying the fingerprinting approach on a sediment core.

Material and methods

SPM sampling strategy



Sample treatment and analysis




Data treatment and geochemical mixing model :

- Kruskal-Wallis test + Discriminant Function Analysis → tracer selection
- Mixing model + Monte Carlo analysis (1000 repetitions) → relative source contribution estimates + uncertainties
- Data treatment : standardisation, particle size correction (Gellis and Noe, 2013)

Residual metal concentrations in SPM samples

Residual Al, Ba and Zn concentrations in SPM in Rhône River tributaries




- Al reflect the grain size
- Ba is a discriminating element but not Zn at the Upper Rhône scale
→ confirmed by Kruskal-Wallis test and DFA

Source fingerprinting in the Upper Rhône

Geochemical mixing model results from SPM data : relative source contributions in %

Low water flow ($Q \leq 800 \text{ m}^3/\text{s}$) - 2012

Flood ($Q > 800 \text{ m}^3/\text{s}$) - 2012




Source contribution (in %, corrected data) with absolute uncertainties at Jons station

	Arve	Fier	Guiers	Ain	Bourbre
Low water flow (n = 13)	95 ± 2	2 ± 2	1 ± 3	1 ± 6	1 ± 4
Flood (n = 6)	60 ± 10	10 ± 11	14 ± 19	9 ± 12	7 ± 16
Dam flushing (n = 8)	95 ± 2	0	3 ± 3	1 ± 2	1 ± 2

Preliminary results on sediment core

Residual and total Al, Ba and Zn concentrations :



These results will be used to trace the history of sediment sources at the Arles stations.

Conclusions

- Original approach to trace with the residual fraction of SPM in Upper Rhône
- Discriminant elements were found
- Robust relative contribution results at the sample and Upper Rhône scales

Perspectives

- Try to reduce uncertainties of relative contributions by applying a particle size correction factor [1]
- Complete the OSR SPM database as a number of samples are missing, on some tributaries, to trace SPM sources at Arles station and in sediment core
- Historical SPM inputs in the Upper Rhône
- Compare SPM and sediment core results