Geophysical Research Abstracts Vol. 21, EGU2019-12926, 2019 EGU General Assembly 2019 © Author(s) 2019. CC Attribution 4.0 license.



Using multi-source satellite data to assess recent snow-cover change and its uncertainty in the Tibet Plateau

Yingsha Jiang, Fei Chen, Yanhong Gao, Michael Barlage, and Jianduo Li Northwest Institute of Eco-environment and Resources, CAS, Key Laboratory of Land Surface Process and Climate Change in Cold and Arid Regions, China (jiangys@lzb.ac.cn)

Snow cover in the Qinghai-Tibet Plateau (QTP) is a critical component in the water cycle and regional climate of East Asia. Satellite data from three different sources (i.e. the Multi-Sensor Synergy of FengYun-3 (FY3A/B/C MULSS), the Moderate Resolution Imaging Spectroradiometer (MODIS), and the Interactive Multi-sensor Snow and Ice Mapping System (IMS)) were used to analyze the QTP fractional-snow-cover (FSC) change and associated uncertainties in the last decade. A four-step cloud removal procedure was applied to those data, which effectively reduced the cloud percentage from 40.8-56.1% to 2.2-3.3%, with an averaged error about 2% estimated by a random sampling method. Compared with in-situ observations, the cloud-removed FY3B and MODIS have annual mean classification accuracy of about 94%, which is higher than other FY3 data and IMS. Five data sets (FY3A/B/C, MODIS, IMS) show significant differences in snow cover fraction, extent, variation trends and duration changes. IMS has the widest snow cover extent with highest FSC (4.7% and 9.5% higher than MODIS and FY3B FSC, respectively). FY3A/C and MODIS, on morning overpass satellites, have around 5% higher FSC than afternoon-overpass FY3B. The annual FSC and snow duration decrease for 2012-2017 shown by at least two of the FY3B, MODIS and IMS data, but they increase for IMS and decreases for MODIS in recent decade (2006-2017) due to different changing rates of snow onset and melting. Meanwhile, their variation rates increase with the increasing elevation bands lower than 6 km. For uncertainties, higher elevation bands (<6 km) have higher uncertainties, and snow-melting trend has higher uncertainty than snow-onset trend.