



Response of stratospheric water vapor and ozone to the unusual timing of El Niño and the QBO disruption in 2015–2016

Mohamadou Diallo (1), Martin Riese (1), Thomas Birner (2), Paul Konopka (1), Rolf Müller (1), Michaela I. Hegglin (3), Michelle L. Santee (4), Mark Baldwin (5), Bernard Legras (6), and Felix Ploeger (1)

(1) Forschungszentrum Juelich, Institute of Energy and Climate Research Stratosphere (IEK-7), Juelich, Germany (m.diallo@fz-juelich.de), (2) Meteorological Institute, Faculty of Physics, Ludwig-Maximilians-University Munich, Germany, (3) Department of Meteorology, University of Reading, Reading, UK, (4) Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA, (5) College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK, (6) Laboratoire de Météorologie Dynamique, UMR8539, IPSL, UPMC/ENS/CNRS/Ecole Polytechnique, Paris, France

The stratospheric circulation determines the transport and lifetime of key trace gases in a changing climate, including water vapor and ozone, which radiatively impact surface climate. The unusually warm El Niño–Southern Oscillation (ENSO) event aligned with a disrupted Quasi-Biennial Oscillation (QBO) caused an unprecedented perturbation to this circulation in 2015–2016. Here, we quantify the impact of the alignment of these two phenomena in 2015–2016 on lower stratospheric water vapor and ozone from satellite observations. We show that the warm ENSO event substantially increased water vapor and decreased ozone in the tropical lower stratosphere. The QBO disruption significantly decreased global lower stratospheric water vapor and tropical ozone from early spring to late autumn. Thus, this QBO disruption reversed the lower stratosphere moistening triggered by the alignment of the warm ENSO event with westerly QBO in early boreal winter. Our results suggest that the interplay of ENSO events and QBO phases will be crucial for the distributions of radiatively active trace gases in a changing future climate, when increasing El Niño-like conditions and a decreasing lower stratospheric QBO amplitude are expected.