



## **Biogeochemical Controls of CO<sub>2</sub> Concentrations and Fluxes in an Amazon Floodplain Lake – a Multiscale Approach**

Joao Henrique Fernandes Amaral (1,2), John Melack (2,3), Pedro Maia Barbosa (2), Alberto Vieira Borges (5), Sally MacIntyre (2,6), Daniele Kasper (1), Alicia Cortes Cortes (6), and Bruce Rider Forsberg (1)

(1) Coordenação de Dinâmica Ambiental, Laboratório de Ecossistemas Aquáticos, Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil, (2) Earth Research Institute, University of California, Santa Barbara, California, (3) Bren School of Environmental Science and Management, University of California, Santa Barbara, California, USA, (5) Chemical Oceanography Unit, University of Liège, Liège, Belgium, (6) Marine Science Institute, University of California, Santa Barbara, California, USA

Seasonal variations in inundation and related environmental conditions in floodplains govern carbon dioxide dynamics. Variations in CO<sub>2</sub> concentrations and exchanges with the atmosphere were investigated over 24 hour periods from August 2014 to September 2016 in two different regions of Lake Janauacá, an Amazon floodplain lake (3°23' S 60°18' W). Meteorological, ancillary environmental measurements, and temperature and dissolved oxygen profiles were combined with pCO<sub>2</sub> in surface waters and chamber estimates of air-water gas exchange. pCO<sub>2</sub> values varied by four orders of magnitude (6 to 66289  $\mu$ atm) and were greater during the night. CO<sub>2</sub> emissions varied by two orders of magnitude (- 5 to 119 mmol m<sup>-2</sup> h<sup>-1</sup>) and, in general, were greater during the day. Wind protected sites, closer to vegetated areas, had higher pCO<sub>2</sub> values and lower CO<sub>2</sub> fluxes compared to sites with more wind exposure. Inter-annual differences were mainly related to changes in herbaceous plant coverage associated with exceptionally low water levels in one year, while diel changes were related to planktonic dynamics. pCO<sub>2</sub> values were positively correlated with water depth and pCO<sub>2</sub> measured in aquatic plant mats, but inversely related to chlorophyll. Our findings demonstrate how variations in inundation govern the different biogeochemical controls of CO<sub>2</sub> dynamics in floodplains, and how these environments can respond to exceptional conditions.