



Spatial-temporal dynamics of dissolved nitrate and its source identification in the upper Han River basin, China

Jie Yuan (1), Hongmei Bu (2), Yanxin Wang (1), and Quanfa Zhang (3)

(1) School of Environmental Studies, China University of Geosciences, Wuhan, 430074, P.R. China, (2) Key Laboratory of Water cycle and Related Land Surface Processes, Institute of Geographic sciences and Natural Resources Research, Chinese Academy of Sciences (CAS), Beijing 100101, P. R. China, (3) Key Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical Garden, CAS, Wuhan 430074, P. R. China

Nitrate (NO_3^-) contamination, as a major form of nitrogen (N) pollution, is a severe environmental problem in river ecosystems with intensive human activities. Source identification of NO_3^- contamination in rivers is pivotal for better management of water quality. Here, we investigated the spatial-temporal dynamics of dissolved NO_3^- in the upper Han River (including the mainstream and major tributaries) with intensive industrial and agricultural disturbance in central China using data from 32 sample sites at four periods, and identified the NO_3^- sources using data of stable nitrogen ($\delta^{15}\text{N}-\text{NO}_3^-$), oxygen and hydrogen ($\delta^{18}\text{O}-\text{H}_2\text{O}$ and $\delta\text{D}-\text{H}_2\text{O}$) isotopes. A great deal of spatial-temporal variation in NO_3^- concentrations was observed with the highest values in summer (22.75 ± 17.75 mg/L) compared with other seasons. Moreover, cluster analysis divided the sampling sites into three clusters, representing low, moderate and high N pollution level, respectively. The $\delta\text{D}-\text{H}_2\text{O}$ and $\delta^{18}\text{O}-\text{H}_2\text{O}$ compositions indicated that modern precipitation was the major water source for the river. Furthermore, a large range of $\delta^{15}\text{N}-\text{NO}_3^-$ isotope values (from -20.25‰ to 31.46‰) were discovered, implying that the NO_3^- could originate from diverse sources but can be mainly derived from urban or domestic sewage and atmospheric deposition. Overall, our results demonstrated distinct spatial-seasonal differences of NO_3^- pollution and sources, and poor control of N into the river. This study provides useful information for mitigating nitrogen pollution and eutrophication as well as formulating watershed management in river ecosystems.