



Highly explosive basaltic eruptions: magma fragmentation induced by rapid crystallisation

Fabio Arzilli (1), Giuseppe La Spina (1), Mike Burton (1), Margherita Polacci (1), Nolwenn Le Gall (2), Margaret Hartley (1), Danilo Di Genova (3), Biao Cai (4), Nghia Vo (5), Emily Bamber (1), Sara Nonni (5), Robert Atwood (5), Ed Llewellyn (6), and Peter Lee (2)

(1) University of Manchester, School of Earth and Environmental Sciences, Manchester, United Kingdom (fabio.arzilli@manchester.ac.uk), (2) Department of Mechanical Engineering, University College London, London, UK, (3) Institute of Non-Metallic Materials, Clausthal University of Technology, Clausthal-Zellerfeld, Germany, (4) School of Metallurgy and Materials, University of Birmingham, Birmingham B15 2TT, UK, (5) Diamond Light Source, Harwell Science and Innovation Campus, Didcot OX11 0DE, UK, (6) Department of Earth Sciences, Durham University, Durham DH1 3LE, UK

The low viscosity of basaltic magmas generally favours effusive and mildly explosive volcanic activity. Highly explosive basaltic eruptions occur less frequently and their eruption mechanism still remains subject to debate [1] [2] [3] [4], with implications for the significant hazard associated with explosive basaltic volcanism. Particularly, highly explosive eruptions require magma fragmentation, yet it is unclear how basaltic magmas can reach the fragmentation threshold [5] [6].

In volcanic conduits, the crystallisation kinetics of an ascending magma are driven by degassing and cooling [7] [8]. So far, the crystallisation kinetics of magmas have been estimated through ex situ crystallization experiments. However, this experimental approach induces underestimation of crystallization kinetics in silicate melts. The crystallization experiments reported in this study were performed in situ at Diamond Light Source (experiment EE12392 at the I12 beamline), Harwell, UK, using basalt from the 2001 Etna eruption as the starting material. We combined a bespoke high-temperature environmental cell with fast synchrotron X-ray microtomography to image the evolution of crystallization in real time. After 4 hours at sub-liquidus conditions (1170 °C and 1150 °C) the system was perturbed through a rapid cooling (0.4 °C/s), inducing a sudden increase of undercooling. Our study reports the first in situ observation of exceptionally rapid plagioclase crystallisation in basaltic magmas. We combine these constraints on crystal kinetics and viscosity evolution with a numerical conduit model to show that exceptionally rapid syn-eruptive crystallisation is the fundamental process required to trigger basaltic magma fragmentation under high strain rates.

[1] Giordano, D. and Dingwell, D. (2003) *Bulletin of Volcanology*, 65, 8-14. [2] Houghton, B.F. and Gonnermann, H.M. (2008) *Chemie der Erde-Geochemistry*, 68, 117-140. [3] Szramek, L.A. (2016) *Journal of Geophysical Research: Solid Earth*, 121, 7119-7136. [4] Moitra, P. et al. (2018) *Earth and Planetary Science Letters*, 500, 97-104. [5] Papale, P. (1999). *Nature*, 397, 425. [6] Gonnermann, H.M. (2015) *Annual Review of Earth and Planetary Sciences*, 43, 431-458. [7] Cashman, K. and Blundy, J. (2000) *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 358, 1487-1513. [8] La Spina, G. et al. (2016) *Nature Communication*, 7, 13402.