



Changes in cloud cover distribution under different types of atmospheric circulation in Europe

Lucian Sfică (), Christoph Beck (), Andrei Niță (), Andreas Philipp (), Mirela Voiculescu (), and Marius-Victor Bîrsan ()

(1) Alexandru Ioan Cuza University of Iași, Faculty of Geology and Geography, Department of Geography, Iași, Romania (sfical@yahoo.com), (2) Augsburg University, Institute of Geography, Augsburg, Germany (christoph.beck@geo.uni-augsburg.de), (3) Romanian Meteorological Administration, Bucharest, Romania (nitaandru@gmail.com), (4) Augsburg University, Institute of Geography, Augsburg, Germany (a.philipp@geo.uni-augsburg.de), (5) Dunarea de Jos University, Faculty of Science and Environment, Galati, Romania (Mirela.Voiculescu@ugal.ro), (6) Romanian Meteorological Administration, Bucharest, Romania (marius.birsan@gmail.com)

The role of cloud cover in the radiation budget of the planet represents a key element in the understanding of the current climate change. For this reason, nowadays, there is a need for a better understanding of changes in the cloud cover which could modulate the amplitude of climate change at regional or continental scale. In Europe, COST733 catalogue offers the opportunity to disentangle the complex link between cloud cover and the atmospheric circulation. In our study we present the main features of this link using daily data for cloud cover and circulation atmospheric type classification.

NCEP, ERA, CFSR, CMSAF and MERRA cloud cover database were compared and used. The 27 atmospheric circulation types were obtained for two domain size in Europe (domain 00 -as defined in cost733 v.2 - and a smaller one restricted to the continent), applying an objectivised classification method (k-means) with 4 input variables which are relevant for cloud development (mean sea level pressure, 500 hPa height, 850-1000 hPa thickness and relative vorticity at 500 hPa). Both types of data are available for the common period: 1979-2013.

The methodological steps imply firstly the identification of the key regions for the variability of the cloud cover at continental scale applying a principal component analysis to the cloud cover data. Secondly, for the key-regions obtained in the first step, those circulation types which are causing positive or negative anomalies in these areas were selected. In the final step, from these circulation types inducing coherent positive/negative anomalies in the cloud cover only those circulation types characterized by a clear linear trend (negative or positive) were kept. In this way we have identified the regions in Europe where the changes in cloud cover are directly connected with changes in atmospheric circulation.

This analysis delivers coherent results regardless of the domain size used. Two major changes in cloud cover in Europe in connection with atmospheric circulation are depicted. The first one is related to a decrease of the cloud cover in the region of the Baltic Sea in connection with an increase of circulation types associated with atmospheric blocking development over central northern part of the continent. The second one indicates an increase of cloud cover in the south-west part of Europe as a result of a decrease in circulation types associated with the development of the Azores High and its north-eastward ridges.

The results are presented at annual and seasonal level. Also, the climatic impact of these changes is explored.