



## **Transient simulation using the iLOVECLIM-CARAIB coupled model equipped with oxygen isotopes and comparison with ice core $\delta^{18}\text{O}_{\text{atm}}$ data**

Thomas Extier (1), Didier Roche (1,2), Louis François (3), and Amaelle Landais (1)

(1) Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France (thomas.extier@lscce.ipsl.fr), (2) Earth and Climate Cluster, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands, (3) Unité de Modélisation du Climat et des Cycles Biogéochimiques, UR SPHERES, University of Liège, Quartier Agora, Bât. B5c, Allée du six Août 19c, 4000 Liège, Belgium

The  $\delta^{18}\text{O}_{\text{atm}}$  (i.e.  $\delta^{18}\text{O}$  of atmospheric  $\text{O}_2$ ) combines past variations of the low latitude water cycle and of the biosphere productivity. Over the last 800 ka, the  $\delta^{18}\text{O}_{\text{atm}}$  measured in EPICA Dome C (EDC) ice core shows orbital and millennial variations which are similar to the low latitude hydrological cycle variations observed in the  $\delta^{18}\text{O}_{\text{calcite}}$  of the East Asian speleothems. The  $\delta^{18}\text{O}_{\text{atm}}$  is also a precious tool for ice core dating and chronology construction. However, the uncertainty associated with the use of this proxy as chronological constraint is high (6 ka) because of the lack of quantitative interpretation of the  $\delta^{18}\text{O}_{\text{atm}}$  including past evolution of oxygen fluxes and associated isotopic fractionation.

This study presents a modelling approach for a quantitative interpretation of the  $\delta^{18}\text{O}_{\text{atm}}$  through the estimation of the oxygen fluxes and fractionation factors over several glacial-interglacial cycles. This approach combines (1) the coupling between the intermediate complexity climate model iLOVECLIM and the vegetation model CARAIB, (2) an estimation of the oxygen fluxes associated with photosynthesis and respiration processes, (3) an estimation of the isotopic composition of the oxygen in leaf water ( $\delta^{18}\text{O}_{\text{leaf}}$ ) and (4) an estimation of the oxygen fractionation during oxygen uptake. This method associates the climatic parameters and water isotopes simulated by iLOVECLIM with the changes in vegetation distribution and terrestrial biosphere productivity obtained by CARAIB for 26 PFTs (Plant Functional Types) instead of the 2 PFTs previously used in iLOVECLIM.

We present here the results of this new coupled model over several glacial-interglacial cycles and discuss the spatial distribution of the vegetation and  $\delta^{18}\text{O}_{\text{leaf}}$ , as well as the carbon and oxygen fluxes linked to the continental biosphere productivity. A focus on the Last Glacial Maximum can also allow a direct comparison with previous “offline” simulations with General Circulation Models. The  $\delta^{18}\text{O}_{\text{leaf}}$  corresponding to the terrestrial contribution of the  $\delta^{18}\text{O}_{\text{atm}}$ , our results can be compared, as a first approximation, to the isotopic  $\delta^{18}\text{O}_{\text{atm}}$  data measured in EDC ice core.