
Oblique convergence, indentation and bivergent tectonics in Taiwan: insights from field observations and analog models equal to the convergence of the convergence of









*Department of Geosciences, National Taiwan University, Taipei, Taiwan. **Bepartment of Earth Sciences, National Central University, Taiwan. **Bepartment of Earth Sciences, National Central University, Taiwan. **Géosciences Montpellier, UMR 5243 CNRS-University, Taiwan. **Bepartment of Earth Sciences, National Central University, Taiwan. **Department of Earth Sciences, National Central University, National Central University, Taiwan. **Department of Earth Sciences, National Central University, National Central Unive





Tectonic evolutionary model -



In Taiwan today, the subduction of the Chinese continental margin under the Philippine Sea plate results in the progressive growth of an active orogenic wedge. It is one of the best places to study the complex relationships that occur between the tectono-metamorphic processes controlling deformation (plate rheology and kinematics) and surface processes (erosion and sedimentation). In the Central Range of Taiwan, foliation and lineation traces outline the geometry and kinematics of deformation in both, the foreland and hinterland of the orogenic wedge. The foliation dip and the strain ellipsoids distribution show the fan shape of a large pop-up structure characterizing the 3D effects of oblique convergence, indentation and roll-back tectonics. On the eastern flank of the mountain range, regionally developed penetrative cleavage dips, isotope data and sedimentary structures demonstrate regional rollback structures. Two mélange units, the Kenting and Lichi mélange are exposed at the south and east of the Central Range respectively. We present the results of 2D and 3D sandbox models designed to investigate the complex deformation characterizing the active Taiwan orogenic wedge. Experiments allow the study of interactions between tectonics and surface processes, accounting for various boundary conditions and parameters such as sedimentation, erosion, basal friction, and décollement level. They account for the development of those mélanges and bivergent structures. We then characte-


rize the exhumation patterns, the rotation of foliation and stretching lineation

and the displacement patterns induced by strain partitioning in the orogen.





extended continental margin


Analog model accounting for surface processes and structural inheritance





Geodynamic setting of the Taiwan orogenic belt





















Field observations











corresponding author:



Cross sections showing the relationships between cleavage and stretching lineations



Two models for the shape and evolution of strain

