

EGU2020-10219

<https://doi.org/10.5194/egusphere-egu2020-10219>

EGU General Assembly 2020

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Comparing the impact for hydrology of the new ERA5 reanalyses dataset over ERA-Interim for 8 hydrological models in 6 catchments using the eWaterCycle community modelling environment.

Rolf Hut¹, Niels Drost², Jerom Aerts¹, Laurene Bouaziz^{1,3}, Willem van Verseveld³, Bert Jagers³, Fedor Baart³, Edwin Sutanudjaja⁴, Lieke Melsen⁵, Andrew Bennett⁶, Louise Arnal⁷, Fabrizio Fenicia⁸, Leonard Santos⁹, Emiliano Gelati¹⁰, Marco Dal Molin⁸, Wouter Knoben¹¹, Shervan Gharari¹², Caitlyn Hall¹³, Eric Hutton¹⁴, Nick van de Giesen¹, and the the eWaterCycle comparison team from the Netherlands eSciencecenter.*

¹Delft University of Technology, The Netherlands (r.w.hut@tudelft.nl)

²Netherlands eScience Center, The Netherlands

³Deltares, The Netherlands

⁴Utrecht University, Utrecht, The Netherlands

⁵Wageningen University, The Netherlands

⁶University of Washington, USA

⁷European Center for Medium range Weather Forecast (ECMWF), UK

⁸CRP Gabriel Lippmann

⁹Swedish Meteorological and Hydrological Institute (SMHI), Sweden

¹⁰Institut Català de Ciències del Clima, Spain

¹¹University of Bristol, UK

¹²University of Saskatchewan, Canada

¹³Arizona State University, USA

¹⁴Community Surface Dynamics Modeling System (CSDMS), USA

*A full list of authors appears at the end of the abstract

The release of the European Centre for Medium-Range Weather Forecasts (ECMWF)'s Re-Analysis 5 (ERA-5) global climate forcing dataset is expected to greatly improve the quality of hydrological modeling. Following this release there is great interest in assessing the improvements of ERA-5 relative to its predecessor ERA-Interim for hydrological modeling and predictions.

In this study we compare streamflow predictions when using ERA-interim vs ERA-5 as forcing data for a suite of hydrological models from different research groups that capture the variation in modelling strategies within the hydrological modelling community. We check whether physically based models, defined as those that do not require additional parameter calibration, would lead to different conclusions in comparison to conceptual models, defined as those that require calibration. Based on the hydrological model structure we expect that conceptual models that need calibration show less difference in predicting discharge (skill) between ERA-5 and ERA-Interim, where-as the physical based (non-calibrated) models most likely will benefit from the

improved accuracy of the ERA-5 input. This assessment will provide the HEPEX community with answers on how the ERA-5 dataset will improve hydrological predictions based on different hydrological modelling concepts.

An additional key objective while conducting this study is compliance to the FAIR principles of data science. To achieve this we held a workshop in Leiden, the Netherlands, where multiple hydrological models were integrated into the eWatercycle II system. eWatercycle II is a hydrological model platform containing a growing number of hydrological models. The platform facilitates research and cohesivity within the hydrological community by providing an Open-Source platform built specifically to advance the state of FAIR and Open Science in Hydrological Modeling. We also use this study to demonstrate the feasibility of eWatercycle II as a platform for FAIR hydrological models.

Preliminary results from this comparison study were presented at the AGU Fall Meeting 2019. Here we will present the full results of the comparison study.

the eWaterCycle comparison team from the Netherlands eSciencecenter.: Ben van Werkhoven Ronald van Haren Yifat Dzigan Jaro Camphuijsen Fakhreh Alidoost Inti Pelupessy Berend Weel Gijs van den Oord Stefan Verhoeven Bouwe Andela Peter Kalverla