

EGU2020-10320

<https://doi.org/10.5194/egusphere-egu2020-10320>

EGU General Assembly 2020

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



Ocean dynamics and climate during a Neoproterozoic snowball Earth and its aftermath as simulated in a coupled Earth system model

Lennart Ramme and Jochem Marotzke

Max Planck Institute for Meteorology, The Ocean in the Earth System, Germany (lennart.ramme@mpimet.mpg.de)

The Neoproterozoic glaciations, referred to as snowball Earth periods, describe the most extreme transition from a very cold climate to a state of strong greenhouse effect. Atmospheric CO₂ concentrations are rising during the snowball, due to the shutdown of oceanic and terrestrial carbon sinks, until a tipping point is reached and a rapid deglaciation sets in. Subsequently, a warm and completely ice-free climate under very high CO₂ concentrations develops. We show first results of simulations using a coupled atmosphere-ocean general circulation model covering the initiation, as well as the melting of the Marinoan snowball Earth (645 – 635 My ago) and the greenhouse climate in its aftermath. CO₂ concentrations are decreased to initiate a global glaciation and then increased again in order to melt the snowball Earth. As soon as a certain CO₂ threshold is reached, sea-ice melts rapidly, reaching a completely ice-free ocean after only one hundred years, in our model without land glaciers. The ocean becomes strongly stratified, because at the surface the freshwater from the sea-ice melt is warming up quickly, whereas the deeper ocean remains cold and salty. Ocean surface currents return to their pre-snowball behavior soon after the melt, but destratification is slow. The largest mixed layer depths of up to 500 m are reached in the mid latitudes of the winter hemisphere. We compare the climate before and after the snowball state and estimate the time needed for destratification.