



Oxygen-to-nitrogen ratios in 1.5-million-year-old ice cores from Allan Hills Blue Ice Areas: implications for the long-term atmospheric oxygen concentrations

Yuzhen Yan^{1,2}, Michael Bender^{1,3}, Edward Brook⁴, Heather Clifford⁵, Preston Kemeny⁶, Andrei Kurbatov⁵, Sean Mackay⁷, Paul Mayewski⁵, Jessica Ng⁸, Jeffrey Severinghaus⁸, and John Higgins¹

¹Department of Geosciences, Princeton University, Princeton, NJ, USA

²Department of Earth, Environmental and Planetary Sciences, Rice University, Houston, TX, USA (yuzhen.yan@rice.edu)

³School of Oceanography, Shanghai Jiao Tong University, Shanghai, China

⁴College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR, USA

⁵Climate Change Institute, University of Maine, Orono, ME, USA

⁶Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

⁷Department of Earth and Environment, Boston University, Boston, MA, USA

⁸Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, USA

Gases preserved in ice cores provide a potential direct archive for atmospheric oxygen. Yet, oxygen-to-nitrogen ratios in ice cores (expressed as $\delta\text{O}_2/\text{N}_2$) are modified by a number of processes related to gas trapping and gas losses in the ice. Such complications have long hindered the use of ice core $\delta\text{O}_2/\text{N}_2$ to derive true atmospheric oxygen concentrations. Recently, a persistent decline in $\delta\text{O}_2/\text{N}_2$, observed in four different ice cores (GISP2, Vostok, Dome F, and EDC), is interpreted to reflect decreasing atmospheric O_2 concentrations over the late Pleistocene (Stolper et al., 2016). The rate of $\delta\text{O}_2/\text{N}_2$ change is -8.4 ± 0.2 ‰/Myr (1σ). Using new measurements made on EDC samples stored at -50 °C and therefore free from gas loss, Extier et al (2018) confirms the decrease in $\delta\text{O}_2/\text{N}_2$ with a slope of -7.0 ± 0.6 ‰/Myr (1σ).

Here, we present new $\delta\text{O}_2/\text{N}_2$ measurements made on 1.5-million-year-old blue ice cores from Allan Hills Blue Ice Areas, East Antarctica. We use argon-to-nitrogen ratios ($\delta\text{Ar}/\text{N}_2$) in the ice to correct for the fractionations during bubble close-off and gas losses. In those processes, $\delta\text{Ar}/\text{N}_2$ is fractionated in a fashion similar to $\delta\text{O}_2/\text{N}_2$ (Huber et al., 2006; Severinghaus and Battle, 2006). Paired $\delta\text{O}_2/\text{N}_2$ - $\delta\text{Ar}/\text{N}_2$ values measured from the same sample were classified into three different time slices: 1.5 Ma (million years old), 950 ka, and 490 ka. Between 950 ka and 490 ka, we observe a decline in $\delta\text{O}_2/\text{N}_2$ similar to that observed in the aforementioned deep ice cores. This observation gives us confidence in the validity of the Allan Hills blue ice $\delta\text{O}_2/\text{N}_2$ records. Between 1.5 Ma and 950 ka, however, there is no statistically significant trend in ice core $\delta\text{O}_2/\text{N}_2$. Our results show a surprising lack of variability from 1.5 to 0.95 Ma; even during the past ~ 0.9 Ma, the rate of decline was very slow.