



Atmospheric CO₂ during the Late Miocene Cooling

Thomas Tanner, José Guitián, Iván Hernández-Almeida, and Heather Stoll
ETH Zurich, Geological Institute, Earth Science, Zurich, Switzerland (tannerth@student.ethz.ch)

Alkenone sea surface temperature records recently observed suggest a substantial long-term and large-magnitude ocean surface cooling during the Late Miocene. At the same time, starting about seven million years ago, both hemispheres on Earth witnessed synchronous cooling and large areas of the continents experienced drying and enhanced seasonality. Coinciding with this climatic shift were significant changes in ecology, including the rise of C₄-photosynthesizing terrestrial plants and the emergence of so-called "vital effects" in oceanic coccolithophores. These changes are collectively hypothesized to be induced by declining atmospheric CO₂. However, the sparse proxy data available for this time interval limits our understanding of the link between these changes and atmospheric greenhouse gas fluctuations and has let people to propose a "climate-CO₂ decoupling".

In this study, the alkenone based pCO₂ proxy is used to reconstruct atmospheric CO₂ for the time interval between 4.5 and 8.5 Ma. Estimations are based on the carbon isotopic fractionation during photosynthesis (ϵ_p) and a new statistical multilinear regression model based on an analysis of culture and sediment data. Past coccolithophore growth rates are reconstructed using foraminiferal isotopic-based proxies, related to water column structure which favour or limit nutrient supply to the photic zone. A thorough sensitivity analysis of modern and past ϵ_p values and its influencing factors in the Southern Ocean yield to a new, high resolution pCO₂ record. Estimated pCO₂ concentrations synchronously decline with the observed long-term cooling (5°C) from 6.8 to 5.9 Ma, periodically decreasing to sufficiently low values of <200 ppm, potentially inducing ephemeral Northern Hemisphere glaciation. CO₂ concentrations during the Late Miocene Cooling Event are thus successfully reproduced in this study and allow a reasonable interpretation of past conditions as has not yet been previously achieved in the relevant literature.