

EGU2020-18591

<https://doi.org/10.5194/egusphere-egu2020-18591>

EGU General Assembly 2020

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



A pan-African high-resolution drought index dataset

Jian Peng¹, Simon Dadson¹, Feyera Hirpa¹, Ellen Dyer¹, Thomas Lees¹, Diego G. Miralles², Sergio M. Vicente-Serrano³, and Chris Funk^{4,5}

¹University of Oxford, School of Geography and the Environment, Oxford, United Kingdom of Great Britain and Northern Ireland (jian.peng@ouce.ox.ac.uk)

²Laboratory of Hydrology and Water Management, Ghent University, Ghent, Belgium

³Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas (IPE-CSIC) Zaragoza, Spain

⁴U.S. Geological Survey, Earth Resources Observation and Science Center, Sioux Falls, South Dakota

⁵Santa Barbara Climate Hazards Center, University of California, USA

Droughts in Africa cause severe impacts such as crop failure, food shortages, famine, epidemics and even mass migration. To minimize the effects of drought on water and food security over Africa, a high-resolution drought dataset is essential to establish robust drought hazard probabilities and to assess drought vulnerability considering a multi- and cross-sectorial perspective that includes crops, hydrological systems, rangeland, and environmental systems. Such assessments are essential for policy makers, their advisors, and other stakeholders to respond to the pressing humanitarian issues caused by these environmental hazards. In this study, a high spatial resolution Standardized Precipitation-Evapotranspiration Index (SPEI) drought dataset is presented to support these assessments. We compute historical SPEI data based on Climate Hazards group InfraRed Precipitation with Station data (CHIRPS) precipitation estimates and Global Land Evaporation Amsterdam Model (GLEAM) potential evaporation estimates. The high resolution SPEI dataset (SPEI-HR) presented here spans from 1981 to 2016 (36 years) with 5 km spatial resolution over the continent of Africa. To facilitate the diagnosis of droughts of different duration, accumulation periods from 1 to 48 months are provided. The quality of the resulting dataset was compared with coarse-resolution SPEI based on Climatic Research Unit (CRU) Time-Series (TS) datasets, and Normalized Difference Vegetation Index (NDVI) calculated from the Global Inventory Monitoring and Modeling System (GIMMS) project, as well as with root zone soil moisture modelled by GLEAM. Agreement between the coarse resolution SPEI from CRU TS (SPEI-CRU) and the developed SPEI-HR provides confidence in the estimation of temporal and spatial variability of droughts in Africa with SPEI-HR. In addition, agreement of SPEI-HR versus NDVI and root zone soil moisture – with average correlation coefficient (R) of 0.54 and 0.77, respectively – further suggests that SPEI-HR can provide valuable information to study drought-related ecological and societal impacts at sub-basin and district scales in Africa. The dataset is archived in Centre for Environmental Data Analysis (CEDA), with link: <http://dx.doi.org/10.5285/bbdfd09a04304158b366777eba0d2aeb> (Peng et al., 2019).

Peng, J.; Dadson, S.; Hirpa, F.; Dyer, E.; Lees, T.; Miralles, D.G.; Vicente-Serrano, S.M.V.-S.; Funk, C. (2019): High resolution Standardized Precipitation Evapotranspiration Index (SPEI) dataset for Africa. Centre for Environmental Data Analysis, 05 August 2019. doi:10.5285/bbdfd09a04304158b366777eba0d2aeb.