

EGU2020-19481

<https://doi.org/10.5194/egusphere-egu2020-19481>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Deciphering the Signal of Arctic Climate Change

Audrey Morley^{1,2,3}, Markus Raitzsch^{1,4}, Jelle Bijma^{4,5}, Szabina Karancz^{4,6}, and Michal Kucera¹

¹MARUM - Centre for Marine Environmental Sciences, University of Bremen, Bremen, Germany (

²National University of Ireland Galway, Geography, Galway, Ireland

³iCRAG Irish Centre for Research in Applied Geosciences, Ireland

⁴Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar-und Meeresforschung, Bremerhaven, Germany

⁵Jacobs University, Bremen, Germany

⁶Royal Netherlands Institute for Sea Research and Utrecht University, The Netherlands

Whether or not Arctic regions remain(ed) a carbon sink or source to the atmosphere during rapidly warming climates (in the past) is a fundamental question with regards to future global warming and ocean acidification. The boron isotopic composition of planktonic foraminiferal shell calcite ($\delta^{11}\text{B}_{\text{CC}}$) can potentially provide valuable information of past seawater pH if information on a second carbonate system parameter, temperature, and salinity is available. However, most applications of palaeoceanographic proxies to the cold polar oceans are limited due to a paucity of calibration data, limited information on the calcification habitat, and secondary effects of the carbonate system on the temperature recorded by Mg/Ca values measured in the dominant Arctic species *Neogloboquadrina pachyderma sinistral* (NPS). Here we present a new Multi-Collector Inductively Coupled Mass Spectrometry (MC-ICPMS) $\delta^{11}\text{B}$ dataset measured on live NPS collected via plankton tows from the Labrador Sea and Baffin Bay. We compare our results with $\delta^{11}\text{B}_{\text{borate}}$ derived from pH measurements, $\delta^{13}\text{C}$ DIC seawater values, temperature and salinity collected at the time and depth the foraminifera calcified. To quantify the control of low carbonate ion concentration on Mg/Ca derived temperatures we measured B/Ca alongside Mg/Ca in the calibration dataset. We are thus able to present a new geochemical correction scheme that can isolate non-thermal controls on the Mg/Ca-temperature relationship for NPS, allowing us for the first time the reconstruction of carbonate system parameters in the Arctic Ocean.