



## Pacific Meridional Overturning Circulation during the Mid-Pliocene Warm Period

Heather L. Ford<sup>1,2</sup>, Natalie Burls<sup>3</sup>, and David Hodell<sup>2</sup>

<sup>1</sup>Queen Mary University of London, School of Geography, London, United Kingdom of Great Britain and Northern Ireland (h.ford@qmul.ac.uk)

<sup>2</sup>Department of Earth Science, University of Cambridge, Cambridge, United Kingdom of Great Britain and Northern Ireland

<sup>3</sup>Atmospheric, Oceanic, and Earth Sciences, George Mason University, Fairfax, United States of America

Today in the North Pacific only intermediate water forms because of a strong halocline, but Pacific Meridional Overturning Circulation (PMOC) may have existed in the past. The mid-Pliocene warm period (3.264-3.025 Ma) is a time of sustained warmth where atmospheric carbon dioxide concentrations were similar to today and the northern hemisphere was relatively ice free – making it a pseudo-analogue for future climate change. North Pacific sedimentological and climate modeling evidence suggests a PMOC formed during this time. To determine the spatial extent of a PMOC during the mid-Pliocene warm period, we constructed a depth transect of sites between 2400 to 3400 m water depth on Shatsky Rise by measuring stable isotopes of *Cibicides wuellerstorfi*. We compare these new results with previously published records and calculate anomalies using the OC3 water column and core-top data products. The  $\delta^{13}\text{C}$  spatial pattern is consistent with a modest PMOC of intermediate depth (core ~2000 m) extending to the equator during the mid-Pliocene warm period. Ventilation of the North Pacific by a PMOC has broad implications for deep ocean carbon storage as the North Pacific contains the oldest, carbon-rich waters today. Future work will include minor and trace element analyses to determine the temperature and carbon characteristics of the PMOC water mass and comparisons with PlioMIP modeling outputs.