



## Greenhouse climate forces expansion of peatlands into inland areas

Zhihui Zhang<sup>1,2</sup>, Chengshan Wang<sup>1,2</sup>, Dawei Lv<sup>1,3</sup>, and Tiantian Wang<sup>1,4</sup>

<sup>1</sup>State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Beijing, 100083, China

<sup>2</sup>School of Earth Sciences and Resources, China University of Geosciences, Beijing, 100083, China

<sup>3</sup>Shandong Provincial Key Laboratory of Depositional Mineralization and Sedimentary Minerals, Shandong University of Science and Technology, Qingdao, Shandong 266590, China

<sup>4</sup>Institute of Earth Sciences, China University of Geosciences, Beijing, 100083, China

As a significant terrestrial carbon reservoir, peatland has great potential to affect the global carbon cycle and global climate. However, our understanding of broad-scale mechanisms that control the long-term global peatland expansion and carbon accumulation rates is still limited. Here we present a new data synthesis of global coal deposit location, thickness changes, carbon concentration, and distribution area changes, along with new carbon pool estimates of global peatlands from Devonian through geological time. By identifying orbital cycles in coal seams, we show that the long-term rate of carbon accumulation (LORCA) in peatland calculated from published data is controlled by  $pO_2$ ,  $pCO_2$ , temperature, precipitation, and total solar irradiance. We use this relationship and latitudinal temperature gradients to reconstruct the equations between LORCA with latitude on different geological time. The results suggest that there are three main sets of high carbon pool and high carbon accumulation rate of global peatlands in Late Paleozoic, Early-Middle Jurassic, and Late-Cretaceous to Early Cenozoic under low tectonic activity and high terrestrial plant diversity background. In addition, we measure the shortest distances between all coal locations and coastlines based on the new Scotese's paleo-Atlas for the past 400 million years, in order to exhibit the extent of peatland expansion into inland. The result shows the Early-Middle Jurassic period has the longest average distance, which is probably due to the high sea level that minimizes the development of peat swamps on coastal areas and facilitated the moisture to move into deeper inland under the Jurassic Greenhouse climate condition. This study highlights that combining comprehensive coal-related database with paleoclimate, tectonics, and evolution of land plants provides insights into the mechanisms of the long-term behavior of the peatland expansion and carbon reservoir through deep time.

Keywords: Greenhouse climate, Peatland expansion, Carbon pool, Carbon accumulation rates, Coal

This study was financially supported by the National Natural Science Foundation of China (grant No. 41888101), the National Key R&D Plan of China (grant No. 2017YFC0601405) and the National Natural Science Foundation of China (grants 41790450, 41772096).

