

EGU2020-6460

<https://doi.org/10.5194/egusphere-egu2020-6460>

EGU General Assembly 2020

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



## Nutrient sources in four forest watersheds in Japan: the contribution of precipitation, weathering and litter fall

Yoshiyuki Inagaki<sup>1</sup> and Kazumichi Fujii<sup>2</sup>

<sup>1</sup>Forestry and Forest Products Research Institute, Shikoku Research Center, Kochi, Japan (yinagaki@affrc.go.jp)

<sup>2</sup>Forestry and Forest Products Research Institute, Ibaraki, Japan (fkazumichi@affrc.go.jp)

Base cation such as calcium and potassium is important nutrient for tree growth in forest ecosystems. Major sources of basic cation to forest soil are precipitation, weathering of parent material and litter fall. Weathering rate of basic cation is estimated from the input from precipitation, runoff from stream and uptake by vegetation. Nutrient cycling of many Japanese forest ecosystems has been studied but information about the contribution of these sources is limited. This study compared three nutrient sources in four forest watersheds in Japan. These are Oyasan (Gunma Pref.), Kamikamo (Kyoto Pref.), Kiryu (Shiga pref.) and Takatori (Kochi Pref.). We calculated nutrient budget of potassium, calcium and magnesium. The rate of rock weathering of base cation ranged from 0.22 to 4.37 kmolc ha<sup>-1</sup> yr<sup>-1</sup> and increased in the order of Kamigamo < Kiryu < Oyasan < Takatori. In potassium cycling, the contribution of litter fall was greater than that of weathering or precipitation. The runoff of calcium and magnesium from soil was greater in Takatori, where the rate of weathering was high. Although the rate of weathering varied substantially among forest watersheds, the annual flux of litter fall was relatively constant. Similar calculation was applied for nitrogen cycling and the source from soil was treated as that from nitrogen accumulated during the past pedogenesis. Nitrogen input from the accumulated soil source was 36 % in Oyasan whereas that in other three watersheds was 0%. Oyasan was considered as a nitrogen-saturated forest and the result of the study suggest the ecosystem relies on different nitrogen source.