

EGU2020-6909

<https://doi.org/10.5194/egusphere-egu2020-6909>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



## Carbonate and chert genesis in the 3.35 Ga old Strelley Pool Formation (Australia): Insights from trace metals and Sm/Nd dating

Sebastian Viehmann<sup>1</sup>, Simon V. Hohl<sup>2</sup>, Nathalie Tepe<sup>3</sup>, Martin Van Kranendonk<sup>4</sup>, Joachim Reitner<sup>5</sup>, Thilo Hofmann<sup>3</sup>, Christian Koeberl<sup>6</sup>, and Patrick Meister<sup>1</sup>

<sup>1</sup>Universität Wien, Dept. of Geodynamics and Sedimentology, Vienna, Austria (sebastian.viehmann@univie.ac.at)

<sup>2</sup>Tongji University Shanghai, State Key Laboratory of Marine Geology, Shanghai, P.R. China

<sup>3</sup>Universität Wien, Dept. of Environmental Geosciences, Vienna, Austria

<sup>4</sup>University of New South Wales, Australian Centre for Astrobiology, Kensington, Australia

<sup>5</sup>Georg-August-Universität Göttingen, Dept. of Geobiology, Göttingen Germany

<sup>6</sup>Universität Wien, Dept. of Lithospheric Research, Vienna, Austria

Stromatolites are laminated, presumably microbial structures, consisting largely of an authigenic precipitate, thus, providing potential geochemical archives of early Earth aqueous environments and their habitability. In this study, we report trace element and Sm/Nd isotope data from Palaeoarchean stromatolites and adjacent cherts of the Strelley Pool Formation (NW Australia), obtained by ICP-MS and TIMS, to test their reliability as archives for palaeo-environmental reconstruction and to understand authigenic mineral formation.

Stromatolitic carbonates plot together with the stratigraphically underlying Marble Bar cherts on a linear Sm-Nd regression line yielding an age of  $3253 \pm 320$  Ma. In contrast, associated crystal-fan carbonates yield  $2718 \pm 220$  Ma, suggesting that their Sm-Nd isotope system was altered after deposition. Geochronological information via Sm-Nd dating of black and white cherts is limited, probably due to a reset of the isotope system during an unknown Paleoproterozoic or younger alteration event.

Carbonates, as well as white cherts, show shale-normalized rare earth element and yttrium patterns ( $REY_{SN}$ ; except for redox-sensitive Ce and Eu) parallel to those of modern seawater, indicating a seawater-derived origin. Positive  $Eu_{SN}$  anomalies (2.1 - 2.4), combined with heterogeneous  $\Delta Nd_{3.35Ga}$  values (-3.2 to +5.8) within alternating stromatolite laminae, support that seawater chemistry was variably affected by both continental weathering and high-temperature hydrothermal fluids contributing elements of both young mafic or older felsic rocks. In contrast, black cherts show non-seawater like  $REY_{SN}$  patterns and significant amounts of elements leached from the surrounding rocks, masking the pristine geochemical composition of ancient seawater. In conclusion, Archaean stromatolites indeed preserve pristine authigenic phases at the mm-scale that contain signatures representative of the water chemistry prevailing in the depositional environment.