

EGU2020-7732

<https://doi.org/10.5194/egusphere-egu2020-7732>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



## Zn isotopes in deep sea corals: a useful palaeoceanographic archive?

Susan Little<sup>1,2</sup>, Tina van De Flierdt<sup>2</sup>, David Wilson<sup>1</sup>, Mark Rehkämper<sup>2</sup>, Jess Adkins<sup>3</sup>, and Laura Robinson<sup>4</sup>

<sup>1</sup>University College London, Department of Earth Sciences, Gower Place, London, WC1E 6BS, UK

<sup>2</sup>Imperial College London, Department of Earth Science and Engineering, London, SW7 2BP, UK

<sup>3</sup>California Institute of Technology, Pasadena, CA91125, USA

<sup>4</sup>University of Bristol, School of Earth Sciences, Bristol, BS8 1RJ, UK

Zinc (Zn) is an important bioessential trace element. Its distribution in the modern oceans reflects a combination of biological uptake, remineralization and the physical ocean circulation. Furthermore, the partitioning behaviour of Zn ( $D_{Zn}$ ) and its isotopes ( $\delta^{66}Zn$ ) in carbonates has been linked to ambient seawater carbonate chemistry [1-3].

Development of Zn isotopes in carbonates as a palaeoceanographic tool has been hampered by the high concentrations of Zn in contaminating material, such as lithogenic or authigenic (e.g. Fe-Mn oxide) phases. However, deep-sea corals are large enough to be subjected to aggressive physical and chemical cleaning, enabling effective removal of contaminating phases. They also have several other advantages over traditional palaeoclimate archives, including the ability to assign precise absolute ages to individual specimens based on uranium-series dating [4].

Here we present Zn/Ca and  $\delta^{66}Zn$  data for a suite of modern and recent (<1000 yr) deep sea corals from six ocean regions spanning the far North Atlantic to the Tasman Sea. We observe what appears to be species-specific Zn partitioning behaviour, but no clear links between  $D_{Zn}$  or coral  $\delta^{66}Zn$  and ambient seawater carbonate chemistry. Overall, there is good agreement between measured or best-guess modern seawater  $\delta^{66}Zn$  and coral aragonite  $\delta^{66}Zn$  values, suggesting that corals of species *Desmophyllum dianthus* and genus *Caryophyllia* do not significantly fractionate Zn isotopes during calcification. Deep sea corals may thus provide a useful archive of the past ocean Zn isotope composition and its spatial variability.

[1] Marchitto T. M., Curry W. B. and Oppo D. W. (2000). *Paleoceanography* 15, 299–306.

[2] van Dijk, I., de Nooijer, L. J., Wolthers, M., & Reichert, G. J. (2017). *Geochimica et Cosmochimica Acta* 197, 263-277.

[3] Mavromatis, V., González, A. G., Dietzel, M., & Schott, J. (2019). *Geochimica et Cosmochimica Acta* 244, 99-112.

[4] Robinson, L. F., Adkins, J. F., Frank, N., & van de Flierdt, T. (2014), *DSR Part II* 99, 184-198.

