

# Magma ascent and eruption forecasting at Deception Island Volcano (Antarctica) evidenced by $\delta D$ and $\delta^{18}O$ variations

Alvarez-Valero, A.M.<sup>(1,2)</sup>, Aulinas, M.<sup>(3)</sup>, Geyer, A.<sup>(4)</sup>, Gisbert, G.<sup>(5)</sup>, Kereszturi, G.<sup>(6)</sup>, Nuñez-Guerrero, E.<sup>(1,2)</sup>, Polo-Sanchez, A.<sup>(1)</sup>, Sumino, H.<sup>(7)</sup>




(1) Department of Geology, University of Salamanca, Spain (aav@usal.es). (2) Laboratory of Stable Isotopes (NUCLEUS), University of Salamanca, Spain (3) Department of Mineralogy, Petrology and Applied Geology, University of Barcelona, Spain. (4) Institute of Earth Sciences Jaume Almera, ICTJA, CSIC, Barcelona, Spain. (5) Institute of Geosciences, CSIC, Madrid, Spain. (6) Volcanic Risk Solutions, School of Agriculture and Environment, Massey University, Palmerston North, New Zealand. (7) Department of Basic Sciences, University of Tokyo, Japan

## Introduction


Geochemistry of volatiles in active volcanoes provides insights into the magmatic processes and evolution at depth, such as magma evolution and degassing, which can be implemented into volcanic hazards assessment.

### Deception Island (Antarctica)



The island is located at the southwestern end of Bransfield Strait, 100 km north of the Antarctic Peninsula. (Modified from Geyer et al. 2019)


Deception Island is one of the most active volcanoes in Antarctica, with more than 20 explosive eruptive events registered over the past two centuries.



Simplified geological map of Deception Island showing the location of the analyzed samples, and distribution of the two main tectonic fault sets (NW-SE and NE-SW). (Modified from Alvarez-Valero et al. 2020).



Volcanic and magmatic evolution has been strongly marked by the development of a collapse caldera occurred around 4000 years ago.

Hydrogen and oxygen isotopic variations in the volatiles trapped in the Deception Island rocks provide essential information on the processes controlling the magmatic evolution and eruption dynamics in this volcanic suite.



The construction of the Island is separated into three evolutionary stages pre-, syn- and post-caldera. Source: Geyer et al. (2019)

Volcanic rocks are tholeiitic, from basalts to trachyandites and rhyolites, and follow an alkalinity-increasing trend at the upper end of the subalkaline field in the Total Alkali vs. Silica diagram




Total Alkali vs. Silica diagram (Le Bas et al., 1986) for the DI samples. Source: Alvarez-Valero et al. (2020)


## Results

$\delta D$ ,  $\delta^{18}O$  and  $H_2O$ (%) contents were analysed in glass and melt inclusions of pre-, syn- and post-caldera samples.

### Glass and melt inclusions




Examples of melt inclusions (arrows)-bearing phenocrysts (a) syn-caldera olivine and (b) pre-caldera plagioclase immersed in a sideromelane + palagonitic glassy matrix. (Modified from Alvarez-Valero et al., 2020).



Size estimation and shape (3D) reconstruction of a melt inclusion within an olivine crystal obtained by micro-computed tomography (micro-CT). (Modified from Alvarez-Valero et al. 2020).


### $\delta D$ , $\delta^{18}O$ and $H_2O$ (%) contents



Isotopic compositions change geographically depending on the proximity to different external water sources. There is also a correlation between isotopes and fault systems in the island

### Degassing and rehydration processes

Our calculations for degassing in the Deception Island samples show that isotopic ratios and water contents of magmas are consistent with pre- to syn-eruptive degassing in either closed-, open- or mixed- systems of magmas with an initial  $\delta D$  value close to  $-55\text{‰}$ . While pyroclasts follow a closed-system degassing model, lavas degas in open-system conditions.



Calculated degassing curves at closed- (red arrow), open- (black arrow) and mixed-system conditions (green dashed arrow), for the pre- (a), syn- (b) and post-caldera (c) samples. Water content variations from 1.5 to 2.5 wt% slightly change the curves slopes. Source: Alvarez-Valero et al. (2020)


The  $\delta D$  and  $\delta^{18}O$  isotopic values of the less degassed magmas point to no pre-eruptive interaction between magma and external water. This implies (i) a deep magma reservoir where infiltration of surficial water is limited, and (ii) the low porosity of the rock samples did not allow low- $\delta^{18}O$  waters to circulate.

## Conclusions

(i) Fast ascent and quenching of most magmas, preserving pre-eruptive magmatic signal of water contents and isotopic ratios, with local post-emplacment modification by rehydration due to glass exposition to seawater, and by meteoric and fumarolic waters.

(ii) A plumbing system(s) variable with time and currently dominated by closed-system degassing leading to explosive eruptions

(iii) control on the interactions of ascending magmas with the surface waters producing hydrovolcanic activity throughout the two main fault systems



Schematic summary of the isotopic variation at the post-caldera stage of Deception Island highlighting some sample examples under closed- vs. open- vs. mixed-system degassing conditions (Modified from Alvarez-Valero et al. 2020).

## References

Alvarez-Valero, A.M. et al. (2020).  $\delta D$  and  $\delta^{18}O$  variations of the magmatic system beneath Deception Island volcano (Antarctica): Implications for magma ascent and eruption forecasting. Chemical Geology 542, 119595. <https://doi.org/10.1016/j.chemgeo.2020.119595>  
Geyer, et al. (2019). Deciphering the evolution of Deception Island's magmatic system. Scientific Reports, 9, 373. <https://doi.org/10.1038/s41598-018-36188-4>  
Kusakabe, M. et al (2009). Noble gas and stable isotope geochemistry of thermal fluids from Deception Island, Antarctica. Antarctic Science 21, 255-267. <https://doi.org/10.1017/S0954102009001783>

## Acknowledgements

Financial support was provided by the Spanish Government (MINECO projects) RECALDEC (CTM2009-05919-E/ANT), PEVOLDEC (CTM2011-13578-E/ANT), POSVOLDEC (CTM2016-79617-P)/(AEI/FEDER, UE), VOLGASDEC (PGC2018-095693-B-I00)/(AEI/FEDER, UE), and USAL-2019 project (Programa Propio - mod. 1B). A.G. is grateful for her Ramón y Cajal contract (RYC-2012-11024).

How to cite it: Alvarez-Valero, A.M., Aulinas, M., Geyer, A., Gisbert, G., Kereszturi, G., Nuñez-Guerrero, E., Polo-Sanchez, A., Sumino, H. (2020). Magma ascent and eruption forecasting at Deception Island Volcano (Antarctica) evidenced by  $\delta D$  and  $\delta^{18}O$  variations. EGU General Assembly 2020, 4-8 May, Vienna, Austria. EGU2020-12700. <https://doi.org/10.5194/egusphere-egu2020-12700>