




Dominica: transcrustal magmatic system and eruptive halogen budgets

T. d'Augustin, H. Balcone-Boissard, G. Boudon, C. Martel, E. Deloule, P. Bürckel

Under review (minor revisions) in Geochemistry, Geophysics, Geosystems




Geological Context




Ignimbrites (Balcone-Boissard et al. 2018)

Plinian eruptions (this study)



Results



Blue circles correspond to ignimbrites from Balcone-Boissard et al. (2018)




Storage conditions




- MagmaSat : most of inclusions : 50 – 150 MPa
- VolatileCalc :
 - Most of inclusions : 50-200 MPa
 - Degassing path (closed system, 0-1% exsol)
 - Storage pressures :
 - Phase equilibrium requires pressures lower than 200 MPa (≤ 7.1 km deep)
 - Storage temperatures :
 - 860 – 890°C based on OPX-CPX, OPX-melt and CPX-melt equilibrium (Putirka 2008)
 - PLAG-melt and MGT-melt thermometer unreliable because of crystallizations histories
- $\text{Cl}-\text{H}_2\text{O}$ (isobares Webster, 1997) :
 - Less precise than $\text{H}_2\text{O}-\text{CO}_2$
 - Consistent results

- Halogen ratios
 - = $f(\text{source})$
 - Preserved during degassing
 - Identical for both plinian eruptions and ignimbrites
- Trace element ratios
 - = $f(\text{source})$
 - Identical for both plinian eruptions and ignimbrites
- Common source of magmas for all eruptions



Degassing budgets



- Normalized degassing budgets of each element are the same order of magnitude from an eruption to another
- S degassing is the same order of magnitude as Br
- F degassing is 1 order of magnitude higher than S and Br
- Cl degassing is 2 orders of magnitude higher than S and Br
- Halogen elements must be taken into account as reactive species emitted in volcanic plumes

- Boudon, G., Balcone-Boissard, H., Solaro, C., & Martel, C. (2017). *Revised chronostratigraphy of recurrent ignimbritic eruptions in Dominica (Lesser Antilles arc): Implications on the behavior of the magma plumbing system*. Journal of Volcanology and Geothermal Research, 343, 135–154.
- Balcone-Boissard, H., Boudon, G., Blundy, J. D., Martel, C., Brooker, R. A., Deloule, E., et al. (2018). *Deep pre-eruptive storage of silicic magmas feeding Plinian and dome-forming eruptions of central and northern Dominica (Lesser Antilles) inferred from volatile contents of melt inclusions*. Contributions to Mineralogy and Petrology, 173(12), 101.