

A new reverse subchron (C33n.1r) in the Campanian: astronomical duration estimate and geomagnetic/chronostratigraphic implications

EGU General Assembly 2020




jaume.dinares@ingv.it

Dinarès-Turell, Jaume¹, Wolfgring, Erik², Wagreich, Michael²


¹Istituto Nazionale di Geofisica e Vulcanologia (INGV), I-00143 Roma, Italy.

²University of Vienna, Department for Geodynamics and Sedimentology, A-1090 Vienna, Austria.




Postalm section

Stacking pattern along the Campanian interval (upper C33n) of the precession-related basic marl-carbonate lithologic couplets (p) defining the ~100 ky eccentricity minima cycles (Emin bundles) as defined by groups of 2-3 consecutive well defined couplets (see Wagreich et al. (2012), Wolfgring et al. (submitted)).



Detail of the high-resolution sampled interval along the lower part of C33n that comprises the PFSsubchron.



Wagreich M, Hohenegger J, Neuhuber S. 2012. Nannofossil biostratigraphy, strontium and carbon isotope stratigraphy, cyclostratigraphy and an astronomically calibrated duration of the Late Campanian Radotrunca calcarata Zone. *Cretaceous Research* 38:80-96. <https://doi.org/10.1016/j.cretres.2012.04.006>

Wolfgring, E., Wagreich, M., Dinarès-Turell, J., Gier, S., Böhm, K., Sames, B., Spötl, C. & Popp, F. (2018). The Santonian-Campanian boundary and the end of the Long Cretaceous Normal Polarity-Chron: Isotope and plankton stratigraphy of a pelagic reference section in the NW Tethys (Austria). *Newslett. Strat.* 51, 4, 445-476, <https://doi.org/10.1127/nos/2018/0392>

Wolfgring, E., Wagreich, M., Hohenegger, J., Böhm, K., Dinarès Turell, J., Gier, S., Sames, B., Spötl, C. and Jin, S.D.(submitted). An integrated multi-proxy study of cyclic pelagic deposits from the north-western Tethys: the Campanian of the Postalm section (Gosau Group, Austria).