

Gullies on Mars and seasonal ices visualised using the Colour and Stereo Surface Imaging System (CaSSIS)

S. J. Conway(1), A. Pommerehne(2), J. Raack(3), M. Philippe(1), A. McEwen(4), N. Thomas(2), G. Cremonese(5) and the CaSSIS Team. (1) Laboratoire de Planétologie et Géodynamique, CNRS UMR6112, Université de Nantes, France (susan.conway@univ-nantes.fr) (2) Physikalisches Institut, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland, (3) Institut für Planetologie, Westfälische Wilhelms-Universität, Münster, Germany, (4) Lunar and Planetary Institute, University of Arizona, USA (5) Osservatorio Astronomico di Padova, INAF, Padova, Italy.

1. INTRODUCTION

Gullies on Mars were first reported in 2000 [1] and were hailed as evidence of recent liquid water flows on Mars. Since that time monitoring of gullies has revealed they are active today at times of year when the martian surface is at its coldest and when CO₂ is condensed on the surface [2,3]. To the first order the distribution of CO₂ frost [4,5] and the geographic locations of gullies [6] match - between latitudes of 30° and 45° on pole-facing steep slopes and from 45° to the poles on slopes with any orientation. Detailed observations of gullies on martian sand dunes hinted at a complex interplay between frosted and non-frosted surfaces [7] and recent gully-deposits have been emplaced atop frosted gully-fans [3]. In order to explore the relationship between surface frosts and gully-activity further we focus on Sisyphi Cavi near the south pole of Mars, where gully-activity has been studied [8] and CaSSIS obtained a dense temporal coverage in 2018.

3. INITIAL RESULTS

In winter frost covers all surfaces and dark spots and fans are observed on the gully deposits and preferentially around the gully channels (Figure 2a). The dark spots seem to concentrate on sandy surfaces; such as the dunes at the bottom of Figure 2a, consistent with previous observations. These dark fans and spots are interpreted to be the surface expression of gas-jets generated by the sublimation of CO₂ underneath a continuous slab of CO₂ ice on the surface [11]. The jets occur when the pressure fractures the slab ice. The top of the slopes are the first to defrost, as the average surface temperature increases towards 200K (Figure 2b), followed by sun facing parts of the alcoves and channels (Figure 2c). Notice in both Figures 2b and 2c that the base of the alcove remains frost-covered. As the surface temperature approaches and exceeds 250K and the surrounding terrain is completely defrosted, the last parts of the gully to remain frost covered are the fans (Figure 2d). We interpret this to be a result of the fans having slightly lower thermal inertia than the surrounding materials. This lower thermal inertia could be because the fans have a lower content of ground ice (i.e. a thicker lag on top of the ice-table), or because of more recent depositional events. It is at this time of year when gullies are active [8]. Hence we infer that for gully activity to occur there needs to be both frosted and defrosted surfaces available to drive vigorous sublimation of the CO₂ ice [3,7]. Finally, once defrosting has almost fully completed and surface temperatures have reached their seasonal maximum of ~270K the only remaining surface frosts are in pole-facing niches at the base of gully-alcoves (Figure 2e).




Figure 1: Map of CaSSIS image footprints over Sisyphi Cavi on a MOLA colour-coded elevation map with a CTX mosaic as a semi-transparent overlay. Gullies are mapped in red and CaSSIS PAN filter footprints up to March 2020 are in black.

2. Approach

We use CaSSIS colour images in order to study the relationship between surface frosts and gullies in Sisyphi Cavi. CaSSIS has four colour filters: BLU, PAN, RED and NIR (centered on 500, 675, 836 and 937 nm respectively; see [9]); where the BLU filter is particularly useful for picking up surface frosts [10]. The timing of surface defrosting is latitude dependent. Although the images we use in this study are not from the same latitude we are able to place them into “defrosting” order by using surface temperature measurements from the Thermal Emission Spectrometer (TES) aboard Mars Global Surveyor. We assume that the surface materials vary little over this constricted region and hence thermal inertia variations are less important than latitude in controlling defrosting-timing at a regional scale. The general trend of surface temperatures with time varies little between different latitudes in this geographically small region.

5. CONCLUSIONS

- Ices and patterns of defrosting are easy to monitor with CaSSIS.
- Gully-alcoves defrost before the fans and gullies defrost later than surrounding terrain – suggests activity is driven by the availability of “hot” sediment to trigger more efficient sublimation.
- Further work will examine whether frost patterns differ where activity occurs.

Acknowledgements: SJC is grateful for the financial support of CNES in support of her CaSSIS work. The authors thank the spacecraft and instrument engineering teams for the successful completion and operation of CaSSIS. CaSSIS is a project of the University of Bern funded through the Swiss Space Office via ESA's PRODEX programme. The instrument hardware development was also supported by the Italian Space Agency (ASI) (ASI-INAF agreement no. I/018/12/0), INAF/ Astronomical Observatory of Padova, and the Space Research Center (CBK) in Warsaw. Support from SGF (Budapest), the University of Arizona (LPL) and NASA are also gratefully acknowledged.

References: [1] M.C. Malin and K.S. Edgett (2000) *Science*, 288, 2330–2335. [2] C.M. Dundas et al. (2012) *Icarus*, 220, 124–143. [3] C.M. Dundas et al. (2017) *GSL Spec. Pub.* 467. [4] M. Vincendon et al. (2010) *Geophys. Res. Lett.*, 37. [5] S. Piqueux et al. (2015) *Dyn. Mars*, 251, 164–180. [6] T.N. Harrison et al. (2015) *Icarus*, 252, 236–254. [7] K. Pasquon et al. (2019) *Icarus*, 329, 296–313. [8] J. Raack et al. (2015) *Icarus*, 251, 226–243. [9] N. Thomas et al. (2017) *Space Sci. Rev.*, 212, 1897–1944. [10] L.L. Tornabene et al. (2018) *Space Sci. Rev.* 214. [11] H.H. Kieffer et al. (2006) *Nature*, 442, 793–796.




Figure 2: Defrosting of gullies in Sisyphi Cavi as seen by CaSSIS. (a) MY34_002397_317_1, Ls 189° at 07:55 and 73°S. (b) MY34_003751_266_0, Ls 257° at 03:23 and 73°S. (c) MY34_003454_281_0 Ls 241° at 12:38 and 70°S. (d) MY34_003464_256_2, Ls 242° at 07:22 and 68°S. (e) MY34_004349_276_2, Ls 288° at 18:01 and 73°S. Central graph: the general trend of TES surface temperatures for latitudes between 65°S and 75°S, where only the “tail” of falling temperatures towards winter differs by latitude, relative positions of the images a-e are shown.