Energetic particle flux variations detected at low altitudes by Space Application of Timepix Radiation Monitor (SATRAM)

St. Gohl<sup>12</sup>, Frantisek Nemec<sup>2</sup>, Benedikt Bergmann<sup>1</sup>, Stanislav Pospisil<sup>1</sup>

<sup>1</sup>Institute of Experimental and Applied Physics, Czech Technical University in Prague <sup>2</sup>Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic

EGU 2020 - 04./08. May 2020



1/26

IEAP. MFF



St. Gohl

# The Timepix detector





- Developed within the Medipix collaboration at CERN [1]
- Features:
  - 256 x 256 pixels
  - 55 µm pixel pitch
  - 14 bits/pixel
  - Minimal threshold: 3.5 keV
- Bump-bonded to readout chip containing in each pixel cell: amplifier, discriminator, digital counter or ADC or Timer
- Single layer particle tracking capability
- Typical sensor materials: Si, GaAs:Cr, CdTe, CZT

E >

ъ

2/26

IEAP. MFF

#### St. Gohl

# Working principle: Modes of operation







Charges, that are created by ionizing particles hitting the sensor, are separated by the applied bias and create signals at the cathode. A clock is running to measure the time a signal stays above certain threshold (ToT). Through calibration the deposited energy can be determined [2]. Another mode is the ToA, measuring the exact time of the particle impact. Modes cannot be used simultaneously in one pixel.

3/26

ъ

# Proba-V





ъ

4/26

IEAP. MFF



- SATRAM attached to Proba-V prior start (red circle) [3, 4, 5]
- Altitude = 820 km
- Inclination = 98.8°
- Sun-synchronous orbit
- in orbit since May 2013

#### St. Gohl

# Space Application of Timepix RAdiation Monitor (SATRAM) - Technical details



- Power consumption: 2.5W
- Total mass: 380g (107×70×55 mm)
- Radiation tolerance 1 Mrad for the sensor, 100 krad for the electronics
- SATRAM is platform technology demonstrator



■ 3 frame exposure times: 20s, 0.2s and 0.002s

5/26

IEAP. MFF

full information in [6]

# Van Allen Belts





Proba-V orbits Earth at the inner edge of the inner belt.

Inner belt:

- Altitude: 1.000 6.000 km
- Electrons: 0.1 7 MeV
- Flux:  $10^8 \text{ cm}^{-2} \text{ s}^{-1}$
- Protons: 5 MeV 400 MeV
- Flux:  $10^4 \text{ cm}^{-2} \text{ s}^{-1}$

#### Outer belt:

Altitude: 13.000 - 60.000 km

イロト 不得 トイヨト イヨト

-

6/26

IEAP. MFF

- Electrons: 0.1 10 MeV
- Flux:  $10^7 \text{ cm}^{-2} \text{ s}^{-1}$

Due to the slight offset of the belts from Earth's geometric center, the inner Van Allen belt makes its closest approach to the surface at the South Atlantic Anomaly.

St. Gohl

# Theoretical spectra in SATRAM orbit





Spectra expected in SATRAM orbit. Both spectra were created with the SPENVIS online tool [7] by using the AP-8-MIN and Ae-8-MIN models [8, 9]. Electrons and protons left of the red line do not penetrate the aluminium cover of the SATRAM device and are therefore not measured.

IEAP, MFF

7/26

ъ



Measured average dose rate for the years 2015 to 2018. The South Atlantic Anomaly (SAA) is clearly visible as well as the higher radiation near the poles.



#### St. Gohl



Example frame of radiation in an area with low dose rate. The small particle track (1 to 3 pixel) are electrons or X-rays. Long thin tracks represent electrons as well. Roundish tracks with 5 to 10 pixels represent protons hitting the sensor perpendicular. The slightly thicker track in the top right is also a proton. The very long long track is minimum ionizing particle (mip).



#### St. Gohl



Example frame for the polar region region. Electrons produce primarily curly tracks which are very thin (1 pixel wide). Lots of them are visible in this frame. The free rectangular area in the bottom left are masked pixels. They became noisy right after the start and are not considered for the analysis.





Example frame for the SAA. The high flux in this region causes a high occupancy in the frame. Lots of proton particles are visible in between (straight tracks 2 to 4 pixels wide). They show the same directionality. The free rectangular area in the bottom left are masked pixels. They became noisy right after the start and are not considered for the analysis.



## Monthly dose rate





Monthly dose rate associated with the respective time of the year. The peaks in the monthly dose rate can be associated with solar events which cause a geomagnetic strom.

Energetic particle flux variations detected at low altitudes by Space Application of Timepix Radiation Monitor (SATRAM)

IEAP, MFF

12/26

# Solar Proton Event (SPE) in September 2017



13/26

IEAP. MFF



# Solar Proton Event (SPE) in September 2017 - 2



14/26

IEAP. MFF



(a) Measured flux from 04.06. to 16.06.2017 for comparison. No SPE or other events were seen during this time.

(b) Measured flux from 04.09. to 16.09.2017.

The picture on the right side shows the increased dose rate for the time of the event compared to a quiet period on the left.

# Strategy for particle identification



IEAP. MFF

Cluster shape [10], cluster height and energy loss are used to distinguish between particle species [11].



St. Gohl





- LET spectrum from data
- Time period: 01.01.2015 to 31.12.2017
- Includes electrons, MIP's, protons and heavier ions
- From simulation: electrons no higher LET than 10 MeV, protons around 100 MeV (black lines)

ъ

16/26

IEAP. MFF

#### St. Gohl





- First region 0 to 10 MeV
- Shows electrons (curly tracks) and mips (long straight tracks)

17/26

IEAP. MFF

#### St. Gohl





- Second region 10 to 100 MeV
- Shows typical proton tracks with different angles

ъ

18/26

IEAP. MFF

#### St. Gohl





- Last region > 100 MeV
- Shows heavy ions which deposit a lot of energy
- some tracks show secondary electrons emitting from the main track
- small tracks represent protons

19/26

IEAP. MFF

#### St. Gohl

80

60

40

n

-20

-40

-60

-80

Latitude [°] 20

-atitude [°]

-150-100

# Separation of electrons and protons



20/26

IEAP. MFF



 $10^{-2}$ 

were selected to reduce overlapping tracks which causes an underestimation  $10^{-1}$  of the flux in the center of the SAA.

0 Longitude [°] (c) Dominated by lons

50 100 150

St. Gohl

# Geomagnetic storms seen by SATRAM





Revisited the data for analysis of dynamic events in the magnetosphere

Geomagnetic storms seen by SATRAM in the years 2015 to 2018

Selection criteria: Dst<-100nT; set as 0-point in the time axis

Data from 1 day prior event and 6 days after

9 storms selected plotted on top of each other

21/26

IEAP. MFF

St. Gohl

Energetic particle flux variations detected at low altitudes by Space Application of Timepix Radiation Monitor (SATRAM)

# Interplanetary shocks seen by SATRAM

Interplanetary Shocks particles orrs 10<sup>4</sup> 10<sup>4</sup> 10<sup>7</sup> 10<sup>7</sup> 10<sup>7</sup> 10<sup>8</sup> 10<sup>9</sup> 10<sup></sup>

Data from 1 day prior event and 6 days after

10 shocks selected and plotted on top of each other

22/26

IEAP. MFF





# Conclusion



23/26

IEAP. MFF

Summary:

- SATRAM is in space for 7 years
- First successful demonstration of a Timepix working in open space
- Measurement of dose rates depnding on orbital position
- Development of a strategy for particle identification
- Separation of electron, proton or ion dominated fluxes
- Start of a new analysis of space weather observed by SATRAM

Future plans:

- Further analysis on space weather events and include data from other spacecraft (e.g. DEMETER, TARANIS)
- Application of machine learning tools for better separation of electrons and protons

## References I



24/26

IEAP. MFF

- X. Llopart et al. 'Timepix, a 65k programmable pixel readout chip for arrival time, energy and/or photon counting measurements'. In: *Nucl. Instrum. Methods A* 581 (2007), pp. 485–494. DOI: https://doi.org/10.1016/j.nima.2007.08.079.
- J. Jakubek. 'Precise energy calibration of pixel detector working in time-over-threshold mode'. In: *Nucl. Instrum. Methods A* 633 (2011), pp. 262-266. DOI: https://doi.org/10.1016/j.nima.2010.06.183.



ESA Mission overview. 2013. URL: http://www.esa.int/Our\_ Activities/Technology/Proba\_Missions/Proba-V\_carrying\_radiation\_detector\_from\_CERN\_to\_space.



SATRAM aboard Proba-V. 2013. URL: http://www.esa.int/ spaceinimages/Images/2013/04/SATRAM\_payload\_aboard\_Proba-V.



B. Sean. 'The mission of Proba-V - ESA Bulletin'. In: *European Space* Agency 153 (2013), pp. 10–21.

ヘロト 人間 ト ヘヨ ト ヘ ヨ ト ー

# References II

C. Granja et al. 'The SATRAM Timepix spacecraft payload in open space on board the Proba-V satellite for wide range radiation monitoring in LEO orbit'. In: *Planet. Space Sci.* 125 (2016), pp. 114–129. DOI: http://dx.doi.org/10.1016/j.pss.2016.03.009.



Belgian Institute for Space Aeronomy - Space Environment Information System (SPENVIS). ver.4.6.9, Nov. 6. 2017. URL: https://www.spenvis.oma.be/.



- J.I. Vette. *The AE-8 trapped electron model environment*. Tech. rep. NSSDC/WDC-A-RS-91-24, 1991.

St. Gohl

D.M. Sawyer and J.I. Vette. 'AP-8 trapped proton environment for solar maximum and solar minimum'. In: *NASA STI/Recon Technical Report N* 77 (Dec. 1976). Provided by the SAO/NASA Astrophysics Data System. URL: http://adsabs.harvard.edu/abs/1976STIN...7718983S.

25/26

IEAP. MFF

## References III

- T. Holy et al. 'Pattern recognition of tracks induced by individual quanta of ionizing radiation in Medipix2 silicon detector'. In: *Nucl. Intrum. Methods A* 591 (2008), pp. 287–290. DOI: https://doi.org/10.1016/j.nima.2008.03.074.



St. Gohl

St. Gohl et al. 'Study of the radiation fields in LEO with the Space Application of Timepix Radiation Monitor (SATRAM)'. In: *Advances in Space Research* 63.5 (2019), pp. 1646–1660. ISSN: 0273-1177. DOI: https://doi.org/10.1016/j.asr.2018.11.016.

26/26

IEAP. MFF