

Combined use of Sentinel SAR and optical data for soil moisture estimation

Giulia Graldi¹, Simone Bignotti¹, Marco Bezzi² and Alfonso Vitti¹


¹ Dept. of Civil, Environmental and Mechanical Engineering, University of Trento, Italy - ² Tera Group, Trento, Italy

Soil Moisture

Key parameter for studying hydrological processes: its value can be observed punctually or *retrieved by remote sensing*.



© Ambrose Carracho



© ESA/AOES Medialab


Application of Interest

Precision agriculture



© 2014CIAT/GeorginaSmith

Weather forecast models



© NASA

Dataset and Study Case

- **Satellite Images:** Sentinel 1 (VV,VH polarization), Sentinel 2
- **Agricultural Mask:** Corine Land Cover 2018
- **Observations:** REMEDHUS soil moisture network (20 sites)

Study Area: 900km² of agricultural fields, Spain

Study Period: 2018 (18 paired images)

Method and Tools



CHANGE DETECTION: Maximum soil moisture range within the study period for each acquisition date [1]

- Negligible roughness variation with 100m resolution
- Elaboration with equal vegetation condition
- Backscatter coefficient variation only due to soil moisture variation



Trend of the maximum σ_0^{VV} variations

Examples of the distribution of the maximum differences of σ_0^{VV} in the study sample in function of two vegetation indices. In the used method the fitting polynomial of the distribution is used in the soil moisture assessment to scale lower σ_0^{VV} differences. The red fitting polynomial is the one proposed in literature, in green the one used in the elaborations.



Vegetation Indices Tested

Optical Indices

$$NDVI = \frac{NIR - RED}{NIR + RED}$$

$$NDWI = \frac{NIR - SWIR}{NIR + SWIR}$$

NIR: near infrared

SWIR: short wavelength infrared

SAR Indices

$$FVI = \frac{\sigma_0^{VV}}{\sigma_0^{VH}}$$



$$RVI = \frac{4\sigma_0^{VH}}{\sigma_0^{VV} + \sigma_0^{VH}}$$

σ_0^{VV} : VV polarized backscatter coeff.


σ_0^{VH} : VH polarized backscatter coeff.

Maps of soil moisture variation


Maps produced with optical indices have a more speckled spatial distribution if compared to the ones produced with SAR indices.




(c) NDVI



(d) NDWI



(e) FVI



(f) RVI

Soil moisture variation on 14-08-2018

Punctual Validation

N° data	272		$NDVI$	$NDWI$	FVI	RVI
$U.M.$	$\frac{m^3}{m^3}$	MAE	0.042	0.065	0.053	0.058
		$bias$	0.035	0.059	0.045	0.048
		$RMSE$	0.073	0.097	0.083	0.088

Conclusions and Future Developments

- Best performance for NDVI followed by both SAR indices. It would be interesting to further investigate the possibility of using only SAR images with a wider dataset, since SAR is not weather dependent.
- More recognisable spatial patterns ascribable to agricultural fields for SAR indices elaborations.

- Increase of the dataset
- Study of the 2 trends of the SAR indices
- Comparison of SAR and optical indices
- Temporal study of the spatial variations
- Elaboration with higher resolution

References and Contacts

[1] Gao, Q. (2017). "Synergetic Use of Sentinel-1 and Sentinel-2 Data for Soil Moisture Mapping at 100 m Resolution". In: Sensors 17, p. 1966.

giulia.graldi@unitn.it
alfonso.vitti@unitn.it
bezzi@tera-group.it