

EGU21-4048, updated on 25 Oct 2021

<https://doi.org/10.5194/egusphere-egu21-4048>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Relative sea level change during MIS 3: a black hole in the world. New observations from Calabria, central Mediterranean sea

Fabrizio Antonioli¹, Lucio Calcagnile², Luigi Ferranti³, Giuseppe Mastronuzzi⁴, Carmelo Monaco⁵, Paolo Montagna⁶, Paolo Orrù⁷, Gianluca Quarta⁸, Fabrizio Pepe⁹, Giovanni Scardino¹⁰, Giovanni Scicchitano¹¹, Paolo Stocchi¹², and Marco Taviani¹³

¹INGV, Associate Res. Roma, Italy (fabrizioantonioli2@gmail.com)

²CEDAD, University of Salento, Italy (lucio.Calcagnile@unisalento.it)

³Dipartimento di Scienze della Terra, dell'Ambiente e delle Risorse, Napoli University, Italy (luigi.ferranti@unina.it)

⁴Dipartimento di Scienze della Terra, Bari University, Italy (gimastronuzzi@libero.it)

⁵Dipartimento di Scienze della Terra, Catania University, Italy (cmonaco@unict.it)

⁶Polar Sciences, CNR Bologna, Italy (paolo.montagna@cnr.it)

⁷Dipartimento di Scienze della Terra, Cagliari University (orrap@unica.it)

⁸CEDAD, University of Salento, Italy (gianluca.quarta@unisalento.it)

⁹Dipartimento di Scienze della Terra, Palermo University (fabrizio.pepe@unipa.it)

¹⁰Dipartimento di Scienze della Terra, Bari University, Italy (giovanni.scardino@uniba.it)

¹¹Dipartimento di Scienze della Terra, Bari University, Italy (gianfranco.scicchitano@uniba.it)

¹²NIOZ Royal Netherlands Institute for Sea Research Coastal Systems Department, The Nederland (paolo.Stocchi@nioz.nl)

¹³CNR Bologna, Italy (marco.taviani@cnr.it)

Estimates of global ice volume during MIS 3 (60-29 ka) can be constrained between -25 and -87 m (Shackleton, 2000; Waelbroeck et al., 2002; Clark et al., 2009; Hughes et al., 2013; Grant et al., 2014). As regards the maximum altitude reached during this period there are few observed data for a comparison between the global curves and the variations due to different rheostay of the mantle in coastal areas. Uncertainties on the rheostatic behaviour near- or far-fields from the ice bulk during cold period, make it very difficult to estimate the local sea level during MIS 3. Several factors make investigations of MIS 3 sea level difficult: i) the areas where suitable coastal sediments formed are currently submerged at depths of few tens of meters below present sea level; ii) the preservation of geomorphic features and sedimentary records is limited due to the erosion occurred during the Last Glacial Maximum (LGM) with sea level at depth of -130m, followed by marine transgression that determined the development of ravinement surfaces).

Few data were observed worldwide, especially when tectonics or GIA in the near field leads to uplifts. Our research aims to point out what has been published globally and in the Mediterranean, but, above all, to illustrate the sections of new outcrops in Cannitello (Calabria, Italy) where we have found and dated fossiliferous marine pocket beaches deposited on uplifted bed metamorphic rock. Radiocarbon ages of marine shells (about 43 kyrs cal BP) indicate that these outcrops (presently at 28 and 30 meters above sea level) belong to MIS 3.1. Based on some

considerations regarding the altitude of MIS 3.1 highstand, the correction for altitude with the local vertical tectonic movements and GIA of the Cannitello outcrops allows us to revise the eustatic altitude of this highstand. This is consistent with the recent findings (Gowan et al., 2020), which are based on a novel ice sheet modelling technique.

Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica, J.X., Hostetler, S.W., McCabe, A.M., 2009. The Last Glacial Maximum. *Science* 325, 710–714. doi:10.1126/science.1172873

Gowan, E.J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. C., Gyllencreutz, R., Mangerud, J., Svendsen, J. I., Lohmann, G. (in print): Global ice sheet reconstruction for the past 80000 years. *PANGAEA, Earth & Environmental Science* <https://doi.org/10.1594/PANGAEA.905800>.

Grant, K.M., Rohling, E.J., Ramsey, C.B., Cheng, H., Edwards, R.L., Florindo, F., Heslop, D., Marra, F., Roberts, A.P., Tamisiea, M.E., Williams, F., 2014. Sea-level variability over five glacial cycles. *Nature Communications* 5, 5076. doi:10.1038/ncomms6076

Hughes, P.D., Gibbard, P.L., Ehlers, J., 2013. Timing of glaciation during the last glacial cycle: evaluating the concept of a global 'Last Glacial Maximum' (LGM). *Earth-Science Reviews* 125, 171–198. doi:10.1016/j.earscirev.2013.07.003

Shackleton, N.J., 2000. The 100,000-Year Ice-Age Cycle Identified and Found to Lag Temperature, Carbon Dioxide, and Orbital Eccentricity. *Science* 289, 1897–1902. doi:10.1126/science.289.5486.1897

Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J.C., McManus, J.F., Lambeck, K., Balbon, E., Labracherie, M., 2002. Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records. *Quaternary Science Reviews, EPILOG* 21, 295–305. doi:10.1016/S0277-3791(01)00101-9