

EGU21-5999

<https://doi.org/10.5194/egusphere-egu21-5999>

EGU General Assembly 2021

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Venus, an Astrobiology Target

Sanjay Limaye¹, Rakesh Mogul², Kevin Baines³, Mark Bullock⁴, Charles Cockell⁵, James Cutts³, Diana Gentry⁶, James Head⁷, Kandis-Lea Jessup⁸, Vladimir Kompanichenko⁹, Yeon Joo Lee¹⁰, Richard Mathies¹¹, Tetyana Milojevic¹², Rosalyn Pertzborn¹, Lynn Rothschild⁶, Dirk Schulze-Makuch¹³, David Smith⁶, and Michael Way¹⁴

¹University of Wisconsin-Madison, Space Science and Engineering Center, Madison, Wisconsin, United States of America (sanjayl@ssec.wisc.edu)

²Cal Poly, Pomona, California, USA

³Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

⁴Science and Technology Corp., Hampton, Virginia, USA

⁵University of Edinburgh, Edinburgh, UK

⁶NASA Ames Research Center, Mountain View, California, USA

⁷Planetary Science Institute, Tucson, Arizona, USA

⁸Brown University, Providence, Rhode Island, USA

⁹Southwest Research Institute, Boulder, Colorado, USA

¹⁰ICARP, Russian Academy of Science, Birobidzhan, Russia

¹¹Technical University of Berlin, Berlin, Germany

¹²University of California, Berkeley, California, USA

¹³University of Vienna, Vienna, Austria

¹⁴Tokyo University of Technology, Tokyo, Japan

The interest in the possibility of life on Venus is driven not just by curiosity about life originating in another Earth-like environment, but because of the possibility that life may be playing a critical role in the planet's present, and possibly its past, atmospheric state. The brilliance of Venus in the night sky (as viewed from Earth) is due to its highly reflective cloud cover, about 28 km thick at the equator. Its spectral albedo is about 90% at wavelengths > 500 nm, but it drops gradually to about 40% around 370 nm before rising slightly at shorter wavelengths. This albedo drop is due to the presence of several absorbers in the atmosphere and the cloud cover. A very large fraction of the energy absorbed by Venus is at ultraviolet wavelengths with sulfur dioxide above the clouds contributing to the absorption below 330 nm; however, the identities of the other absorbers remain unknown. The inability to identify the absorbers that are responsible for determining the radiative energy balance of Venus over the last century is a major impediment to understanding how the planet "works", a major component of NASA's efforts in planetary exploration. Limaye et al. (*Astrobiology* **18**, 1181-1198, 2018) presented a hypothesis suggesting that cloud-based microbial life could be contributors to the spectral signatures of Venus' clouds, building upon previous suggestions of the possibility of life in the clouds of Venus.

Four interconnected themes for the exploration of Venus as an astrobiology target are: - (i)

investigations focused on the likelihood that liquid water existed on the surface in the past leading to the potential for the origin and evolution of life, (ii) investigations into the potential for habitable zones within Venus' clouds and Venus-like atmospheres, (iii) theoretical investigations into how active aerobiology may impact the radiative energy balance of Venus' clouds and Venus-like atmospheres, and (iv) application of these investigative themes towards better understanding the atmospheric dynamics and habitability of exoplanets. These themes can serve as a basis for proposed Venus Astrobiology Objectives and suggestions for measurements for future missions, as per the goals and objectives developed by the Venus Exploration Analysis Group (VEXAG), which is sponsored by NASA to plan for the future exploration of Venus.

A Venus Collection to be published in *Astrobiology* journal in 2021 will include papers from the "Habitability of the Venus Cloud Layer", Moscow (October 2019) workshop.