

EGU21-6168

<https://doi.org/10.5194/egusphere-egu21-6168>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Daily cyclicality in bivalve shell chemistry: Paleo-weather record or circadian rhythm?

Niels de Winter^{1,2}, Lukas Fröhlich³, Daniel Killam⁴, Wim Boer⁵, Lennart de Nooijer⁵, Gert-Jan Reichart⁵, and Bernd Schöne³

¹Utrecht University, Geosciences Faculty, Earth Sciences, Utrecht, Netherlands

²AMGC research group, Vrije Universiteit Brussel (VUB), Brussels, Belgium

³Institute of Geosciences, University of Mainz, Mainz, Germany

⁴Biosphere 2, University of Arizona, Oracle, Arizona, United States

⁵Department of Ocean Systems, Royal Netherlands Institute for Sea Research (NIOZ), Den Burg, Texel, the Netherlands

Bivalve shells have a long-standing reputation as archives for high-resolution (seasonal scale) (paleo)climate variability due to their incremental growth, yielding accurate shell chronologies, and their abundance, diversity, and high preservation potential in the fossil record (Schöne and Surge, 2012). Capitalizing on innovations in geochemical techniques, high-resolution sclerochronology can now resolve changes in bivalve shell chemistry beyond the daily resolution (e.g. Sano et al., 2012; Warter et al., 2018). When applied on fossil shells, these ultra-high-resolution records have the potential to bridge the gap between climate and weather reconstructions and yield unprecedented information about bivalve paleobiology, extreme weather events in past climates and even astronomical cycles (Warter and Müller, 2017; de Winter et al., 2020; Yan et al., 2020).

However, studies of sub-daily scale shell chemistry are almost exclusively limited to giant clams (*Tridacna* spp.), due to their high growth rates. It is hitherto unknown if and how such diurnal cycles in chemistry differ in other genera across the bivalve clade and/or whether they are exclusive to photosymbiotic clams. In addition, it is not clear whether the daily cycles are formed in response to environmental conditions (e.g. light or temperature sensitivity) or reflect circadian rhythms.

To answer these questions, we combine ultra-high-resolution (hourly scale) Laser Ablation ICP-MS trace element profiles through shells of various tridacnid species from the tropical Gulf of Aqaba with profiles through the giant scallop (*Pecten maximus*) from the temperate Atlantic coast of northwestern France. We observe trace element cycles on in the daily frequency domain in both tridacnids and pectinids. This shows that these diurnal cycles are formed regardless of shell mineralogy (aragonite vs. calcite), living environment (tropical inter-tidal vs. temperate sub-tidal) and occur in highly unrelated bivalve taxa. Our data helps the interpretation of similar records from fossil shells in terms of past (extreme) weather events, climate, and shell growth.

References

- de Winter, N. J. et al. *Paleoceanography and Paleoclimatology* 35, e2019PA003723 (2020).
- Sano, Y. et al. *Nature Communications* 3, 761 (2012).
- Schöne, B. R. & Surge, D. M. *Treatise Online* 24, Volume 1, Chapter 14 (2012).
- Warter, V., Erez, J. & Müller, W. *Palaeogeography, Palaeoclimatology, Palaeoecology* 496, 32–47 (2018).
- Warter, V. & Müller, W. *Palaeogeography, Palaeoclimatology, Palaeoecology* 465, 362–375 (2017).
- Yan, H. et al. *PNAS* 117, 7038–7043 (2020).