

EGU21-7247

<https://doi.org/10.5194/egusphere-egu21-7247>

EGU General Assembly 2021

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Sclerochronological evidence of pronounced seasonality from the Pliocene of the southern North Sea Basin, and its implication

Andrew Johnson¹, Annemarie Valentine², Melanie Leng³, Bernd Schöne⁴, Hilary Sloane³, and Stijn Goolaerts⁵

¹School of Environmental Sciences, University of Derby, Derby DE22 1GB, United Kingdom (a.l.a.johnson@derby.ac.uk)

²School of Geography and Environmental Science, Nottingham Trent University, Southwell NG25 0QF, United Kingdom (drmitzyvalentine@gmail.com)

³National Environmental Isotope Facility, British Geological Survey, Keyworth NG12 5GG, United Kingdom (mjl@bgs.ac.uk)

⁴Institute of Geosciences, University of Mainz, 55128 Mainz, Germany (schoeneb@uni-mainz.de)

⁵Directorate Earth and History of Life, Royal Belgian Institute of Natural Sciences, 1000 Brussels, Belgium (sgoolaerts@naturalsciences.be)

Various elements of the biota of the early Pliocene Coralline Crag Formation (southern North Sea Basin, eastern England) have been taken to indicate a warm temperate marine climate, with summer surface temperatures above 20 °C and winter temperatures above 10 °C [1]. However, summer and winter temperature estimates from oxygen-isotope ($\delta^{18}\text{O}$) sclerochronology of benthic invertebrates are typically in the respective cool temperate range when calculated using a plausible modelled value for water $\delta^{18}\text{O}$ of +0.1‰. For instance, examples of the bivalve mollusc *Aequipecten opercularis* from the Ramsholt Member indicate summer maximum temperatures of 11.0–15.7 °C and winter minimum temperatures of 4.4–7.1 °C [2]. Amongst other evidence, the pattern of microgrowth-increment variation in Ramsholt-Member *A. opercularis* points to a depth below the summer thermocline, hence the temperatures recorded for that season provide an underestimate of surface temperature; this may well have been in the warm temperate summer range [2], as suggested by the pelagic dinoflagellate biota [3]. However, the cool temperate benthic winter temperatures indicated by isotopic data are likely also to have obtained at the surface, pointing to a greater seasonal range in surface temperature (perhaps > 15 °C) than in the modern North Sea (< 13 °C) [2]. This conclusion is not changed by adoption of a different (invariant) value for water $\delta^{18}\text{O}$ and also follows from data for a specific late Pliocene interval (Mid-Piacenzian Warm Period) elsewhere in the southern North Sea Basin (Belgium, Netherlands [4]). Here we present isotopic evidence of a seasonal range in surface temperature higher than now at other times in the late Pliocene. Examples of *A. opercularis* from several horizons in the Lillo Formation (Belgium) and the Oosterhout Formation (Netherlands) indicate seasonal ranges in benthic temperature of 10–14 °C. Seasonal variation in water $\delta^{18}\text{O}$ can only plausibly account for about 1 °C of these ranges. Taking into consideration microgrowth-increment evidence of a setting below the summer thermocline, the seafloor ranges imply that the surface seasonal range was sometimes 17 °C or more. Other bivalves (*Atrina fragilis*, *Arctica islandica*, *Pygocardia rustica*, *Glycymeris radiolyrata*) do not indicate such a high seasonal range in benthic (and hence

surface) temperature but this can be attributed to inadequate sampling—time-averaging or a failure to recover evidence of seasonal extremes because of growth breaks. The high surface temperature range could reflect a reduction in vigour of the North Atlantic Current and hence diminished oceanic supply of heat in winter.

References:

- [1] Vignols et al. (2019), *Chem. Geol.* 526, 62–83. <https://doi.org/10.1016/j.chemgeo.2018.05.034>.
- [2] Johnson et al. (2020), *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 561. <https://doi.org/10.1016/j.palaeo.2020.110046>.
- [3] Head (1997), *J. Paleontol.* 71, 165–193. <https://doi.org/10.1017/S0022336000039123>.
- [4] Valentine et al. (2011), *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 309, 9–16. <https://doi.org/10.1016/j.palaeo.2011.05.015>.