

EGU21-9128

<https://doi.org/10.5194/egusphere-egu21-9128>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



## Characteristics and variability of ocean ventilation in the high-latitude North Atlantic in an eddy-permitting ocean model

Helen L. Johnson<sup>1</sup>, Graeme MacGilchrist<sup>2</sup>, David P. Marshall<sup>3</sup>, Camille Lique<sup>4</sup>, Matthew Thomas<sup>5,6</sup>, Laura Jackson<sup>7</sup>, and Richard Wood<sup>7</sup>

<sup>1</sup>Department of Earth Sciences, University of Oxford, Oxford, UK

<sup>2</sup>Atmospheric and Oceanic Science, Princeton University, Princeton, New Jersey, USA

<sup>3</sup>Department of Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, UK

<sup>4</sup>Ifremer, Laboratoire d'Océanographie Physique et Spatiale, Brest, France

<sup>5</sup>National Oceanic and Atmospheric Administration/Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey, USA

<sup>6</sup>University Corporation for Atmospheric Research, Boulder, Colorado, USA

<sup>7</sup>Hadley Centre, Met Office, Exeter, UK

A substantial fraction of the deep ocean is ventilated in the high latitude North Atlantic. As a result, the region plays a crucial role in transient climate change through the uptake of carbon dioxide and heat. We investigate the nature of ventilation in the high latitude North Atlantic in an eddy-permitting numerical ocean circulation model using a set of comprehensive Lagrangian trajectory experiments. Backwards-in-time trajectories from a model-defined 'North Atlantic Deep Water' (NADW) reveal the times and locations of subduction from the surface mixed layer at high temporal and spatial resolution. The major fraction ( $\approx 60\%$ ) of NADW ventilation results from subduction directly into the Labrador Sea boundary current, with a smaller fraction ( $\approx 25\%$ ) arising from open ocean deep convection in the Labrador Sea. There is a notable absence of ventilation arising from subduction in the Greenland-Iceland-Norwegian Seas, due to the re-entrainment of those waters as they move southward. Temporal variability in ventilation arises both from changes in subduction — driven by large-scale atmospheric forcing — and from year-to-year changes in the subsurface retention of newly subducted water, the result of an inter-annual equivalent of Stommel's mixed layer demon. This interannual demon operates most effectively in the open ocean where newly subducted water is slow to escape its region of subduction. Thus, while subduction in the boundary current dominates NADW ventilation, processes in the open ocean set the variability, mediating the translation of inter-annual variations in atmospheric forcing to the ocean interior.