

EGU22-3496

<https://doi.org/10.5194/egusphere-egu22-3496>

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



The analysis of the climate mitigation potential in terms of O₃-Radiative Forcing from aviation NO_x using O₃ algorithmic climate change functions

Pratik Rao¹, Feijia Yin¹, Volker Grewe^{1,2}, Hiroshi Yamashita², Patrick Jöckel², Sigrun Matthes², Mariano Mertens², and Christine Frömming²

¹Delft University of Technology, Delft, Netherlands (p.v.rao@tudelft.nl)

²Deutsches Zentrum für Luft- und Raumfahrt, Institute of Atmospheric Physics, Oberpfaffenhofen, Germany.

Aviation contributes to 3.5% of anthropogenic climate change in terms of Effective Radiative Forcing (ERF) and 5% in terms of temperature change. Aviation climate impact is expected to increase rapidly due to the growth of air transport sector in most regions of the world and the effects of the COVID-19 pandemic are expected to only have a temporary effect on this growth. While efforts have been made to curb CO₂ emissions, non-CO₂ effects that are at least equally significant according to recent research, require more attention. The EU Horizon 2020 project ClimOp considers a comprehensive approach to tackling the climate impact of aviation using novel operational measures. One such measure is climate-optimised flight planning, where small deviations can be made in aircraft trajectories to minimise their overall climate impact. Algorithmic Climate Change Functions (aCCFs) are used to estimate the climate impact of local non-CO₂ effects such as nitrogen oxide (NO_x) emissions (via ozone (O₃) formation and methane (CH₄) depletion), aviation water vapour (H₂O) and contrails using weather variables directly as inputs. By using these functions in an air traffic optimisation module, climate sensitive regions are detected and avoided leading to climate-optimised trajectories. Here, we focus specifically on evaluating the effectiveness of reducing the aviation NO_x induced climate impact via O₃ formation, using only O₃ aCCFs for the optimisation strategy. This is achieved using the chemistry climate model EMAC (ECHAM5/MESSy) and various submodels. A summer and winter day, characterised by high spatial variability of O₃ aCCFs are selected, following which, air traffic over the European airspace is optimised with respect to climate as well as operating cost. The air traffic is laterally and vertically optimised separately to enable an evaluation of the horizontal and vertical pattern of O₃ aCCFs. It is shown that despite the significant impact of the synoptic situation on the transport of emitted NO_x, the climate-optimised flights lead to lower O₃ Radiative Forcing (RF) compared to the cost-optimised flights. The study finds that while O₃ aCCFs can reduce the climate impact, there are certain discrepancies in the prediction of O₃ impact from aviation NO_x emissions, as seen for the selected summer day. Although the aCCFs concept is a rough simplification in predicting future pathways of emissions and subsequent climate impact, we could show that it enables a reasonable first estimate. Further research is required to better describe the aCCFs allowing an improved estimate in O₃-RF reduction for optimisation approaches.

