

EGU23-1353, updated on 29 Sep 2023

<https://doi.org/10.5194/egusphere-egu23-1353>

EGU General Assembly 2023

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



A data-driven approach to derive spatially explicit dynamic "thresholds" for shallow landslide occurrence in South Tyrol (Italy)

Stefan Steger¹, Mateo Moreno^{1,2}, Alice Crespi¹, Stefano Luigi Gariano³, Maria Teresa Brunetti³, Massimo Melillo³, Silvia Peruccacci³, Francesco Marra⁴, Marco Borga⁵, Lotte de Vugt⁶, Thomas Zieher⁷, Martin Rutzinger⁶, Volkmar Mair⁸, Piero Campalani¹, and Massimiliano Pittore¹

¹Institute for Earth Observation, Eurac Research, Viale Druso 1, 39100 Bolzano, Italy (stefan.steger@eurac.edu)

²Faculty of Geo-information Science and Earth Observation (ITC), University of Twente, Hengelosestraat 99, 7514 Enschede, The Netherlands

³Research Institute for Geo-Hydrological Protection, National Research Council (CNR IRPI), Perugia, Italy

⁴Institute of Atmospheric Sciences and Climate, National Research Council (CNR-ISAC), Bologna, Italy

⁵Department of Land, Environment, Agriculture and Forestry, University of Padova, Padova, Italy

⁶Department of Geography, University of Innsbruck, Innrain 52f, 6020 Innsbruck, Austria

⁷Austrian Research Centre for Forests, Rennweg 1, 6020 Innsbruck

⁸Office for Geology and Building Materials Testing, Autonomous Province of Bolzano-South Tyrol, Cardano, Italy

When and where shallow landslides occur depends on an interplay of predisposing, preparatory, and triggering factors. At a regional scale, data-driven analyses are extensively used to assess landslide susceptibility based on "static" maps of predisposing conditions. In contrast, data-driven analyses focusing on landslide triggering factors often rely on non-spatially explicit approaches to derive empirical rainfall thresholds. So far, few attempts have been made to integrate the spatial and temporal analysis domains beyond a posterior combination of separately derived susceptibility models and rainfall thresholds.

This work focuses on the mountainous Italian province of South Tyrol (7400 km²) and proposes a novel data-driven landslide prediction model that jointly considers landslide predisposition and dynamic preparatory and triggering factors. The approach builds on a hierarchical generalized additive model, multi-temporal shallow landslide data from 2000 to 2020 and a range of environmental variables (e.g., daily rainfall, topography, lithology, forest cover). The model produces maps that portray the relative probability of landslide occurrence. These spatially explicit predictions change dynamically as a function of local predisposition, seasonality, and observed (or hypothesized) dynamic preparatory and triggering rainfall (i.e. cumulative rainfall amounts based on varying day-windows). Linking the model output to known measures of model performance, such as hit rate and false alarm rate, enables the creation of dynamic classified maps that can be interpreted in analogy to commonly used empirical rainfall thresholds. The approach also accounts for potential spatial and temporal biases in the landslide inventory by restricting the underlying data sampling to effectively surveyed areas and time periods and by including (and averaging out) bias-describing random effect variables. Our validation confirms the model's high

generalizability and predictive power while providing insights into the interplay of predisposing, preparatory and triggering factors for shallow landslide occurrence in South Tyrol. Application possibilities of this novel approach are discussed.

The research leading to these results is related to the PROSLIDE project that received funding from the research program Research Südtirol/Alto Adige 2019 of the Autonomous Province of Bozen/Bolzano – Südtirol/Alto Adige.