

EGU24-11544, updated on 20 May 2024 https://doi.org/10.5194/egusphere-egu24-11544 EGU General Assembly 2024 © Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Abundance of giant mineral dust particles: Insights from measured emitted dust size distributions during the J-WADI campaign

Hannah Meyer¹, Andres Alastuey², Sylvain Dupont³, Vicken Etyemezian⁴, Jessica Girdwood^{5,6}, Cristina González-Flórez⁷, Adolfo González-Romero^{2,7,8}, Tareq Hussein^{9,10}, Mark Irvine³, Konrad Kandler¹¹, Peter Knippertz¹, Ottmar Möhler¹², George Nikolich⁴, Xavier Querol², Chris Stopford⁵, Franziska Vogel^{12,13}, Frederik Weis¹⁴, Andreas Wieser¹, Carlos Pérez García-Pando^{7,15}, and Martina Klose¹

¹Karlsruhe Institute of Technology (KIT), Karlsruhe, Institute of Meteorology and Climate Research – Troposphere Research (IMKTRO), Germany.

²Institute of Environmental Assessment and Water Research – Consejo Superior de Investigaciones Científicas (IDAEA-CSIC), Barcelona, Spain.

³INRAE, Bordeaux Sciences Agro, ISPA, F-33140 Villenave d'Ornon, France.

⁴Desert Research Institute (DRI), Las Vegas, NV, USA.

⁵University of Hertfordshire, Hertfordshire, UK.

⁶Now at: University of Manchester, Manchester, UK.

⁷Barcelona Supercomputing Center (BSC), Barcelona, Spain.

⁸Polytechnical University of Catalonia (UPC), Barcelona, Spain.

⁹University of Jordan, Amman, Jordan.

¹⁰University of Helsinki, Helsinki, Finland.

¹¹Technical University of Darmstadt, Darmstadt, Germany.

¹²Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate Research – Atmospheric Aerosol Research (IMKAAF), Karlsruhe, Germany.

¹³Now at: Institute of Atmospheric Sciences and Climate (ISAC), National Research Council (CNR), Bologna, Italy.

¹⁴Palas GmbH, Karlsruhe, Germany.

¹⁵Catalan Institution for Research and Advanced Studies (ICREA), Spain.

Gaining a precise understanding of the particle size distribution (PSD) of mineral dust at emission is critical to assess its climate impacts. Despite its importance, comprehensive measurements at dust sources remain scarce and usually neglect part of the super-coarse (particle diameter *d* between 10 and 62.5 μ m) and the entire giant (*d* > 62.5 μ m) particle size ranges. Measurements in those size ranges are particularly challenging due to expected relatively low number concentrations and low sampling efficiencies of instrument inlets.

This study aims to better constrain the abundance of super-coarse and giant dust at emission as part of the Jordan Wind erosion And Dust Investigation (J-WADI, https://www.imk-tro.kit.edu/11800.php) field campaign conducted north of Wadi Rum in Jordan in September 2022. The goal of J-WADI is to improve our fundamental understanding of the emission of desert dust, in particular its full-range size distribution and mineralogical composition.

To capture the dust PSD across the entire size spectrum, we deployed multiple aerosol spectrometers, including active, passive, and open-path devices, such that in combination, a size range from approximately 0.4 to 200 μ m was covered. Here we investigate the variability of the PSD in the super-coarse and giant ranges from observed dust events, address instrumental uncertainties and the impact of different inlets on the resulting PSDs. Our preliminary results reveal a mass concentration peak at around 30 μ m, potentially limited toward larger sizes by substantially reduced inlet efficiencies. Giant dust particles were generally detected during active dust emission starting from friction velocities larger than around 0.2 m s⁻¹.

Based on our results, we will investigate the mechanisms facilitating super-coarse and giant dust particle emission and transport. Quantifying the conditions for and the amount of super-coarse and giant dust at emission will lay the foundation to incorporate its impacts in weather and climate models.