

EGU24-13143, updated on 20 May 2024

<https://doi.org/10.5194/egusphere-egu24-13143>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Land-Atmosphere Coupling Simulation and Its Role in Subseasonal-to-Seasonal Prediction

Yuna Lim^{1,2}, Andrea Molod¹, Randal Koster¹, and Joseph Santanello³

¹Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, Maryland, United States of America

²University of Maryland College Park, College Park, Maryland, United States of America

³Hydrological Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, United States of America

Land-atmosphere (L-A) coupling can significantly contribute to subseasonal-to-seasonal (S2S) prediction. During periods of strong L-A coupling, land-atmosphere feedbacks are expected to enhance the memory of the system and therefore also the predictability and prediction skill. This study aims to evaluate S2S prediction of ambient surface air temperature under conditions of strong versus weak L-A coupling in forecasts produced with NASA's state-of-the-art GEOS S2S forecast system. Utilizing three L-A coupling metrics that together capture the connection between the soil and the free atmosphere, enhanced prediction skill for surface air temperature is observed for 3-4 week boreal summer forecasts across the eastern Great Plains when strong L-A coupling is detected at this lead by all three indices. The forecasts with strong L-A coupling in these "hot spot" regions exhibit warm and dry anomalies, signals that are well simulated in the model. Overall, this study provides insight into how better capturing relevant L-A coupling processes might improve prediction on subseasonal-to-seasonal timescales.