

EGU24-18260, updated on 20 May 2024 https://doi.org/10.5194/egusphere-egu24-18260 EGU General Assembly 2024 © Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Improving daily-to-seasonal sea ice forecasts of the AWI coupled prediction system with sea-ice and ocean data assimilation and atmospheric large-scale wind nudging.

Svetlana Loza¹, Marylou Athanase¹, Longjiang Mu², Jan Streffing¹, Antonio Sánchez-Benítez¹, Miguel Andrés-Martínez¹, Lars Nerger¹, Tido Semmler¹, Dmitry Sidorenko¹, and Helge Goessling¹ ¹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Climate System, Bremerhaven, Germany (svetlana.losa@awi.de)

²Laoshan Laboratory, Qingdao, China

Predictive skills of coupled sea-ice/ocean and atmosphere models are limited by the chaotic nature of the atmosphere. Assimilation of observational information on ocean hydrography and sea ice allows to obtain a coupled-system state that provides a basis for subseasonal-to-seasonal ocean and sea-ice forecast (Mu et al., 2022). However, if the atmosphere is not additionally constrained, the quasi-random atmospheric states within an ensemble forecast lead to a fast divergence of the ocean and sea-ice states, degrading the system's performance with respect to the sea ice forecasts. As reported previously, imposing an additional constraint by nudging largescale winds to the ERA5 reanalysis data (Sánchez-Benítez et al., 2021; Athanase et al., 2022) improves predictive skills of the AWI Coupled Prediction System (AWI-CPS, Mu et al. 2022) with regard to sea ice drift (Losa et al., 2023). Here we provide results based on a much more extensive set of ensemble-based data assimilation experiments spanning the time period from 2002 to 2023 and a series of long forecast experiments over 2010 – 2023, initialized in four different seasons. We compare the performance of forecasts initialized from two sets of data assimilation experiments, with and without atmospheric wind nudging. The additional relaxation of the largescale atmospheric circulation to the ERA5 reanalysis data for the initialization leads to reasonable atmospheric forecast skill on weather timescales: Despite the simple technique, the coarse resolution compared to NWP systems, and the limited optimization efforts, 10-day forecasts of the 500 hPa geopotential height are about as skillful as the best performing NWP forecasts were about 10 –15 years ago. Among other aspects, this leads to significantly improved subseasonal-toseasonal sea-ice concentration and thickness forecasts.

Athanase, M., Schwager, M., Streffing, J., Andrés-Martínez, M., Loza, S., and Goessling, H.: Impact of the atmospheric circulation on the Arctic snow cover and ice thickness variability , EGU General Assembly 2022, Vienna, Austria, 23–27 May 2022, EGU22-5836, https://doi.org/10.5194/egusphere-egu22-5836, 2022.

Losa, S. N., Mu, L., Athanase, M., Streffing, J., Andrés-Martínez, M., Nerger, L., Semmler, T., Sidorenko, D., and Goessling, H. F.: Combining sea-ice and ocean data assimilation with nudging atmospheric circulation in the AWI Coupled Prediction System, EGU General Assembly 2023, Vienna, Austria, 24–28 Apr 2023, EGU23-14227, https://doi.org/10.5194/egusphere-egu23-14227, 2023.

Mu, L., Nerger, L., Streffing, J., Tang, Q., Niraula, B., Zampieri, L., Loza, S. N. and Goessling, H. F. (2022): Sea lice Forecasts With an Upgraded AWI Coupled Prediction System, Journal of Advances in Modeling Earth Systems, 14 (12). doi: 10.1029/2022ms003176

Sánchez-Benítez, A., Goessling, H., Pithan, F., Semmler, T. and Jung, T. (2022): The July 2019 European Heat Wave in a Warmer Climate: Storyline Scenarios with a Coupled Model Using Spectral Nudging, Journal of Climate, 35 (8), pp. 2373-2390. doi: 10.1175/JCLI-D-21-0573.1