

EGU24-2545, updated on 20 May 2024

<https://doi.org/10.5194/egusphere-egu24-2545>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



Copper nanoparticles: Insecticidal action, resistance management and effect on endosymbiont abundance in olive fruit fly *Bactrocera oleae*

Anastasios Malandrakis¹, Kyriaki Varikou², Nektarios Kavroulakis², Antonis Nikolakakis², Irene Dervisi³, Chrysavgi Reppa³, Stefanos Papadakis⁴, Maria Holeva³, and Constantinos Chrysikopoulos⁵

¹Technical University of Crete, School of Chemical and Environmental Engineering, Chania, Greece (tasmal@aue.gr)

²Hellenic Agricultural Organization "ELGO-Dimitra", Institute for Olive Tree, Subtropical Plants and Viticulture, Agrokippio-Souda, 73164 Chania, Greece.

³Benaki Phytopathological Institute, Scientific Directorate of Phytopathology, Laboratory of Bacteriology, Kifissia, 14561, Greece

⁴Department of Biology, University of Crete, Heraklion, 70013, Greece

⁵Department of Civil Infrastructure and Environmental Engineering, Khalifa University of Science and Technology, Abu Dhabi 127788, UAE.

The effectiveness of copper containing nanoparticles (Cu/CuO-NPs) against insecticide-resistant olive fruit flies (*Bactrocera oleae*) and their impact on the insect's reproductive and endosymbiotic parameters were evaluated. The insecticidal activity of both nano and bulk copper [Cu(OH)₂] was comparable or greater than that of the reference insecticide deltamethrin at recommended doses as revealed by feeding experiments. A significant synergistic effect between Cu-NPs or CuO-NPs and deltamethrin was observed in terms of adult mortality. Furthermore, the deltamethrin + Cu-NPs combination decreased the total number of offspring as compared with the untreated control. The above combination also significantly decreased the mean number of stings, pupae, female and total number offspring of the surviving female, compared to deltamethrin applied alone. The abundance of the *Candidatus Erwinia dacicola*- a *B. oleae* larvae bacterial gut endosymbiont- was adversely affected by bulk and nanosized copper. Concluding, Cu-NPs have a great potential to control insecticide-resistant *B. oleae* populations by reducing adult and larval survival and fecundity, and provide the means for reducing the environmental footprint of pesticides by minimizing their required doses.

This study was co-financed by the European Regional Development Fund of the European Union and Greek national funds through the Rural Development Program (RDP/ΠΑΑ) 2014 – 2020, under the call "Cooperation for environmental projects, environmental practices and actions for climate change" (project code: M16SYN2-00354).